

L^∞ instability of Prandtl layers

Emmanuel Grenier* Toan T. Nguyen[†]

October 10, 2019

Abstract

In 1904, Prandtl introduced his famous boundary layer in order to describe the behavior of solutions of incompressible Navier Stokes equations near a boundary as the viscosity goes to 0. His Ansatz was that the solution of Navier Stokes equations can be described as a solution of Euler equations, plus a boundary layer corrector, plus a vanishing error term in L^∞ in the inviscid limit. In this paper we prove that, for a class of smooth solutions of Navier Stokes equations, namely for shear layer profiles which are unstable for Rayleigh equations, this Ansatz is false if we consider solutions with Sobolev regularity, in strong contrast with the analytic case, pioneered by R.E. Caflisch and M. Sammartino [22, 23].

Meanwhile we address the classical problem of the nonlinear stability of shear layers near a boundary and prove that if a shear flow is spectrally unstable for Euler equations, then it is non linearly unstable for the Navier Stokes equations provided the viscosity is small enough.

1 Introduction

In this paper we address the question of the description of solutions of incompressible Navier Stokes equations in a bounded domain, in the case of the zero Dirichlet boundary condition. More precisely, let Ω be the half

*Equipe Projet Inria NUMED, INRIA Rhône Alpes, Unité de Mathématiques Pures et Appliquées., UMR 5669, CNRS et École Normale Supérieure de Lyon, 46, allée d'Italie, 69364 Lyon Cedex 07, France. Email: Emmanuel.Grenier@ens-lyon.fr

[†]Department of Mathematics, Penn State University, State College, PA 16803. Email: nguyen@math.psu.edu. TN's research was partly supported by the NSF under grant DMS-1764119 and by an AMS Centennial Fellowship.

plane $x \in \mathbb{R}$, $y > 0$. Let u^ν be solutions of incompressible Navier Stokes equations with forcing term f^ν

$$\partial_t u^\nu + (u^\nu \cdot \nabla) u^\nu - \nu \Delta u^\nu + \nabla p^\nu = f^\nu, \quad (1.1)$$

$$\nabla \cdot u^\nu = 0 \quad (1.2)$$

and Dirichlet boundary condition

$$u^\nu = 0 \quad \text{on} \quad y = 0. \quad (1.3)$$

As the viscosity goes to 0, we expect u^ν to converge to a solution of Euler equations

$$\partial_t u^E + (u^E \cdot \nabla) u^E + \nabla p^E = f^0, \quad (1.4)$$

$$\nabla \cdot u^E = 0 \quad (1.5)$$

with boundary condition

$$u_2^E = 0 \quad \text{on} \quad y = 0. \quad (1.6)$$

The justification of this convergence is however very delicate, since the boundary conditions dramatically change. As a consequence, a boundary layer is expected near $y = 0$ in order to describe the transition between Navier Stokes boundary conditions and Euler boundary conditions. To take into account this transition, Prandtl [21] introduced the following Ansatz

$$u^\nu(t, x, y) = u^E(t, x, y) + u^P(t, x, y/\sqrt{\nu}) + o(1)_{L^\infty}, \quad (1.7)$$

where u^P describes the behavior of u^ν in a boundary layer of size $\mathcal{O}(\sqrt{\nu})$, called the Prandtl's boundary layer, and the remainder $o(1)_{L^\infty}$ tends to zero in the inviscid limit. The boundary layer corrector $u_1 = u_1^E(t, x, 0) + u_1^P(t, x, Y)$ is then constructed by solving the classical Prandtl boundary layer equation

$$\begin{aligned} \partial_t u_1 + u_1 \partial_x u_1 + u_2 \partial_Y u_1 &= \partial_Y^2 u_1 - \partial_x p^E(t, x, 0) \\ \partial_x u_1 + \partial_Y u_2 &= 0 \end{aligned} \quad (1.8)$$

together with the no-slip boundary conditions $u_1 = u_2 = 0$ at $Y = 0$ and the matching condition $u_1(t, x, Y) \rightarrow u_1^E(t, x, 0)$ as $Y \rightarrow \infty$.

The existence and uniqueness of solutions to the Prandtl equations have been constructed for monotonic data by Oleinik [19] in the sixties. There are also recent reconstructions [1, 18] of Oleinik's solutions via a more direct energy method. For data with analytic or Gevrey regularity, the well-posedness

of the Prandtl equations is established in [22, 5], among others. In the case of non-monotonic data with Sobolev regularity, the Prandtl boundary layer equations are known to be ill-posed ([3, 6, 14]).

Concerning the validity of Prandtl's Ansatz (1.7), this was established for data with analytic regularity in the celebrated work of Caffisch and Sammartino [23]. In particular, it was proven that if a boundary layer Ansatz exists to describe the limiting behavior of u^ν , then it must be of the Prandtl's form (1.7). A similar result were also obtained by [16] for data whose initial vorticity is compactly supported away from the boundary. The stability of shear flows under perturbations with Gevrey regularity is recently proved in [4].

However, considering analytic or Gevrey initial data is too restrictive, since it precludes small but high frequencies perturbations, which are more physically relevant. The first author proved in [7] that the Ansatz (1.7) is nonlinearly unstable with a vanishing lower bound of order $\mathcal{O}(\nu^{1/4})$ for the remainder. Up to now, there were no result which proved, or disproved, the Ansatz (1.7) for data with Sobolev regularity.

In this paper we give the first result in this direction. Namely we prove that there exists particular initial data such that (1.7) is *wrong*. More precisely we will show that some shear layer profiles are nonlinearly unstable, for these profiles the remainder in (1.7) reaches order one in the inviscid limit. Proving the instability of order one, or in fact any order beyond $\mathcal{O}(\nu^{1/4})$ obtained in [7], faces a serious obstruction: the viscous boundary sublayers which arise from the instability of the main Prandtl's layer are themselves unstable, giving rise to thinner and thinner viscous sublayers. The instability of these thinner sublayers is inevitable due to the linear instability theory of generic shear flows [2, 9] and the fact that the local Reynolds is of order $\frac{U_{sub}}{\nu^{1/4}} \rightarrow \infty$ whenever the amplitude of sublayers U_{sub} goes beyond $\nu^{1/4}$. As a consequence, there are many instabilities from both the main Prandtl's layers and the sublayers, and it remains unclear which sublayers are dominant in the large time. This is the main limitation of the previous method [7]. For more details of the obstruction, see Section 2. See also [11] for a further link between the stability of classical Prandtl's layers and that of viscous sublayers.

This paper not only proves the invalidity of the Ansatz (1.7), but also constructs a three-layer solution to Navier-Stokes equations involving an Euler flow (trivial), a classical Prandtl's layer with thickness of order $\sqrt{\nu}$, and a thinner boundary sublayer with thickness of order $\nu^{3/4}$. This latter sublayer in turn gives rise to thinner sub-sublayers with thickness of order

$\nu^{7/8}$, which is confirmed linearly [9]. This paper builds the first step towards fully justifying the boundary layer cascade developed near the boundary.

Let us mention that if one replaces the classical no-slip boundary condition (1.3) by a Navier-slip condition, the boundary layers are less violent with a much smaller amplitude of order $\sqrt{\nu}$. As a consequence, the inviscid limit and the boundary layer Ansatz are established in this case: see for instance [15, 17]. The instability observed in this paper does not apply to these settings. However, when the slip length is of order $\sqrt{\nu}$ or smaller, a similar instability up to order one can be obtained via an energy method [20], adapted from [7]. A crucial difference between the slip and no-slip boundary conditions is that there are intricate (and unstable) boundary sublayers arising in the latter case, but not in the former.

1.1 Main results

Let us now detail our main results of this paper. A shear layer profile is a solution of the form

$$U^\nu(t, x, y) = \begin{pmatrix} U(t, y/\sqrt{\nu}) \\ 0 \end{pmatrix}$$

that is a solution of both Prandtl and Navier Stokes equations. Here $U(t, y)$ is a smooth function with $U(t, 0) = 0$ such that $U(t, y)$ converges when $y \rightarrow +\infty$ to a constant Euler flow U_∞ . In this paper we consider two cases

- time dependent boundary layers:

$$\partial_t U^\nu - \partial_{yy} U^\nu = 0,$$

namely U^ν is a solution of the classical heat equation

- time independent boundary layers: in this case, we add a time-independent forcing term which compensates for the viscosity. Precisely, we take

$$F^\nu = \begin{pmatrix} -U''(0, y/\sqrt{\nu}) \\ 0 \end{pmatrix}. \quad (1.9)$$

The main result of this paper is as follows.

Theorem 1.1. *There exists a smooth, analytic function $U(0, Y)$, such that the corresponding sequence of time dependent shear layers*

$$U^\nu(t, x, y) = \begin{pmatrix} U(t, y/\sqrt{\nu}) \\ 0 \end{pmatrix}, \quad (1.10)$$

which are smooth solutions of Navier Stokes, Prandtl, and heat equations satisfies the following assertion. For any N and s arbitrarily large, there exist $\sigma_0 > 0$, $C_0 > 0$ and a sequence of solutions u^ν of Navier Stokes equations (1.1)-(1.3) with forcing terms f^ν , on some interval $[0, T^\nu]$, such that

$$\|u^\nu(0) - U^\nu(0)\|_{H^s} \leq \nu^N,$$

$$\|f^\nu\|_{L^\infty([0, T^\nu], H^s)} \leq \nu^N,$$

but

$$\|u^\nu(T^\nu) - U^\nu(T^\nu)\|_{L^\infty} \geq \sigma_0$$

and

$$T^\nu = O(\sqrt{\nu} \log \nu^{-1}).$$

This theorem proves that Prandtl Ansatz is false in L^∞ in very small times, of order $\sqrt{\nu} \log \nu^{-1}$. The same theorem holds true for the time independent boundary layer with a forcing term (1.9). In particular, it is proved that the convergence of Navier-Stokes solutions to Euler solutions, plus a boundary layer, fails in L^∞ in the inviscid limit. We remark that this however does not prevent the convergence to hold in L^p for $p < \infty$.

In addition, as will be clear from the construction, the Navier-Stokes solutions obtained in Theorem 1.1 involve not only the Prandtl's layer of size $\sqrt{\nu}$, but also a viscous sublayer of size $\nu^{3/4}$. Moreover, it is important to note that the instability occurs in the vanishing time T^ν of order $\sqrt{\nu} \log \nu^{-1}$.

In the proof of Theorem 1.1, we introduce the following hyperbolic rescaling

$$(T, X, Z) = \frac{1}{\sqrt{\nu}}(t, x, z).$$

Theorem 1.1 is thus a direct consequence of the scaling and the following theorem, which also addresses the classical stability problem of shear layers in the inviscid limit.

Theorem 1.2. *Let $U(y)$ be a smooth and analytic function which converges exponentially fast at infinity to a constant, with $U(0) = 0$ and assume that it is spectrally unstable for linearized Euler equations, with a simple eigenvalue. Then it is nonlinearly unstable for Navier Stokes equations in L^∞ , provided ν is small enough, in the following sense. For any s arbitrarily large, there exist $\sigma_0 > 0$, $C_0 > 0$ and a sequence of solutions u^δ of Navier Stokes equations (1.1)-(1.3) with forcing terms f^δ , on some interval $[0, T^\delta]$, such that, as $\delta \rightarrow 0$,*

$$\|u^\delta(0) - U^\nu(0)\|_{H^s} \leq \delta,$$

$$\|f^\delta\|_{L^\infty([0, T^\nu], H^s)} \leq \delta,$$

but

$$\|u^\delta(T^\delta) - U^\nu(T^\delta)\|_{L^p} \geq \sigma_0, \quad \forall p \in [1, \infty]$$

and

$$T^\delta = O(\log \delta^{-1}),$$

where $U^\nu(t, y)$ is the solution of heat equation with diffusivity ν and initial data $U(y)$.

Let us now discuss this result. Up to the best of our knowledge it is the first rigorous result of instability of a shear layer profile near a boundary for Navier Stokes equations. According to Rayleigh's criterium the profile U^ν will have an inflection point. Physically, this may correspond to a reverse flow and thus rules out the exponential profile $U_\infty(1 - e^{-y/C})$. The Prandtl equation is well posed for U^ν and for neighboring analytic profiles. However we do not know whether the Prandtl equation is well posed for nearby profiles with only Sobolev regularity.

Notations

For $\alpha \in \mathbb{R}$ we define

$$\Delta_\alpha = \partial_y^2 - \alpha^2$$

and

$$\nabla_\alpha = (i\alpha, \partial_y).$$

In particular $\nabla_\alpha^2 = (-\alpha^2, \partial_y^2)$. The three dimensional case is exactly similar to the two dimensional one, therefore we restrict ourselves to the two dimensional case.

2 General strategy

The proof of Theorem 1.1 relies on the *complete* construction of the instability u^ν . The first step is to make an isotropic change of variables in t , x and y ; namely, we define

$$T = \frac{t}{\sqrt{\nu}}, \quad X = \frac{x}{\sqrt{\nu}}, \quad Y = \frac{y}{\sqrt{\nu}}.$$

Of course, the Navier Stokes equations remain unchanged, except the viscosity which is now $\sqrt{\nu}$. From now on, we abuse the notation by denoting by t , x and y the new variables T , X and Y .

The starting point is the choice of the shear layer profile (1.10). We will choose a shear profile $U_0 = (U(y), 0)$ which is unstable with respect to linearized Euler equations. More precisely, we start from U_0 , such that there exists an exponentially growing solution to the following linearized Euler equations

$$\partial_t v + (U_0 \cdot \nabla)v + (v \cdot \nabla)U_0 + \nabla p = 0, \quad (2.1)$$

$$\nabla \cdot v = 0, \quad (2.2)$$

$$v_2 = 0 \quad \text{on} \quad y = 0. \quad (2.3)$$

The study of the linear stability of a shear layer profile is a classical issue in fluid mechanics. The classical strategy to address this question is to introduce the stream function of v and to take its Fourier transform in the tangential variable x (with dual Fourier variable α) and the Laplace transform in time (with dual variable $\lambda = -i\alpha c$). Precisely, we look for v of the form

$$v = \nabla^\perp \left(e^{i\alpha(x-ct)} \psi(y) \right) + \text{complex conjugate}. \quad (2.4)$$

Putting (2.4) in (2.1), we get the classical Rayleigh equation for the stream function ψ

$$(U - c)(\partial_y^2 - \alpha^2)\psi = U''\psi, \quad (2.5)$$

$$\alpha\psi(0) = \lim_{y \rightarrow +\infty} \psi(y) = 0. \quad (2.6)$$

The study of the linear stability of U reduces to a spectral problem: find c and ψ , solutions of Rayleigh equations, with $\Im(\alpha c) > 0$. Following the classical Rayleigh criterium, if such an instability exists, then U must have an inflection point. Such smooth unstable profiles do exist (see, for instance, [7]). We choose the most unstable mode, namely the largest $|\alpha \Im c|$. Starting with such an instability, we can construct an instability for the following linearized Navier Stokes equations

$$\partial_t v + (U_0 \cdot \nabla)v + (v \cdot \nabla)U_0 - \sqrt{\nu} \Delta v + \nabla p = 0, \quad (2.7)$$

$$\nabla \cdot v = 0, \quad (2.8)$$

$$v = 0 \quad \text{on} \quad y = 0. \quad (2.9)$$

The analogs of the Rayleigh equations (2.5)-(2.6) are the Orr Sommerfeld equations which read

$$-\varepsilon(\partial_y^2 - \alpha^2)^2 \psi + (U - c)(\partial_y^2 - \alpha^2)\psi = U''\psi, \quad (2.10)$$

$$\alpha\psi(0) = \psi'(0) = \lim_{y \rightarrow +\infty} \psi(y) = 0, \quad (2.11)$$

where

$$\varepsilon = \frac{\sqrt{\nu}}{i\alpha}.$$

Such a spectral formulation of the linearized Navier-Stokes equations near a boundary layer shear profile has been intensively studied in the physical literature. We in particular refer to [2, 24] for the major works of Heisenberg, Tollmien, C.C. Lin, and Schlichting on the subject. We also refer to [8, 9, 10] for the rigorous spectral analysis of the Orr-Sommerfeld equations.

Now starting from an unstable mode (ψ^0, c^0) of the Rayleigh equation for some positive α , it is possible to construct an unstable mode (ψ^ν, c^ν) for Navier Stokes equations, provided ν is small enough, such that

$$\psi^\nu - \psi^0 = O(\nu^{1/4}), \quad (2.12)$$

$$c^\nu - c^0 = O(\nu^{1/4}). \quad (2.13)$$

Let

$$\lambda_0 = \alpha c^\nu.$$

This has been proved rigorously in [10] through a complete analysis of the Green function of Orr Sommerfeld equation. More precisely, the proof of (2.12)-(2.13) relies on the complete description of all four independent solutions of the fourth order differential equation (2.10). It can be proven that two of them go to $+\infty$ as $y \rightarrow +\infty$. These two solutions can be forgotten in the construction of an unstable mode. The other two converge to 0 as $y \rightarrow +\infty$. One, called ψ_f , has a "fast" behavior, namely behaves like $\exp(-Cy/\sqrt{\varepsilon})$ for large y . The other one, called ψ_s , has a "slow" behavior and behaves like $\exp(-C|\alpha|y)$. The second one, ψ_s , comes from the Rayleigh mode, and is a small perturbation of ψ^0 . Then the unstable mode ψ^ν is a combination of ψ_f and ψ_s and is of the form

$$\psi^\nu = \alpha^\nu \psi_f + \beta^\nu \psi_s. \quad (2.14)$$

The two relations $\psi^\nu(0) = \partial_y \psi^\nu(0) = 0$ give the dispersion relation. Using the fact that ψ_s is an approximate eigenmode for Rayleigh equation, (2.12) and (2.13) can be proved using an implicit function theorem (see [10] for complete details). We also get, for every positive k , that

$$|\partial_y^k \psi^\nu(y)| \leq \frac{C_k}{|\varepsilon^{(k-1)/2}|} e^{-Cy/|\sqrt{\varepsilon}|} + C_k e^{-\beta y} \quad (2.15)$$

for some positive β . Note that ψ^ν has a boundary layer behavior. This is natural since there is a change of boundary conditions between Rayleigh and Orr Sommerfeld equations. The size of the boundary layer is of order $\varepsilon^{1/2} \approx \nu^{1/4}$ (for fixed α), which is introduced to balance $\varepsilon \partial_y^4$ and ∂_y^2 . This sublayer is known as "viscous sublayer" in the physical literature [2]. Note that ψ_s and ψ_f are analytic on a strip $|\Im y| \leq \sigma_0$ for some $\sigma_0 > 0$.

Once the linear instability is constructed, we may construct an approximate solution of the form

$$u^{app}(t, x, y) = \sum_{j=1}^M \nu^{Nj} u^j(t, x, y) \quad (2.16)$$

starting from the maximal unstable eigenmode

$$u^1(t, x, y) = \Re\left(\psi^\nu e^{i\alpha(x-c^\nu t)}\right).$$

The construction of such an approximate solution is routine work, and involves successive resolutions of linearized Navier Stokes equations

$$\begin{aligned} \partial_t u^n + (U_0 \cdot \nabla) u^n + (u^n \cdot \nabla) U_0 - \sqrt{\nu} \Delta u^n + \nabla p^n &= F_n, \\ \nabla \cdot u^n &= 0, \end{aligned} \quad (2.17)$$

together with the zero initial data and zero Dirichlet boundary conditions, with

$$F_n = - \sum_{1 \leq j \leq n-1} (u^j \cdot \nabla) u^{n-j}.$$

Note that by construction [7], u^{app} solves Navier Stokes equations, up to a very small term R^M , of order $\nu^{N(M+1)} e^{(M+1)\Re \lambda_\nu t}$, with $\lambda_\nu = -i\alpha c^\nu$, the maximal unstable eigenvalue.

Let u^ν be the solution of Navier Stokes equations with initial data $u^{app}(0)$. A natural next step is to try to bound the difference $v := u^\nu - u^{app}$ in L^2 norm. However, we only get

$$\frac{d}{dt} \|v\|_{L^2}^2 \leq C(1 + \|\nabla u^{app}\|_{L^\infty}) \|v\|_{L^2}^2 + \|R^M\|_{L^2}^2.$$

As there is a boundary layer in u^{app} , $\|\nabla u^{app}\|_{L^\infty}$ is unbounded as $\nu \rightarrow 0$, and thus, this energy inequality is useless when $u^{app} - U$ is of order greater than $\nu^{1/4}$. Using only energy estimates, we cannot obtain $O(1)$ instability in L^∞ , and are limited to $O(\nu^{1/4})$ instability (this is the main limitation of [7]).

The reason of this failure is that the viscous sublayer becomes linearly unstable in the inviscid limit [2, 9]. The next natural idea is to work with analytic initial data and to hope that analyticity will kill sublayer instabilities, exactly as in Caffisch and Sammartino work [23], where the authors used analyticity to kill any instability of Prandtl's layers. However in the current setting, we want to get control over time intervals of order $\log \nu^{-1}$, namely on unbounded time intervals. As the analyticity radius decreases with time, it becomes small, of order $1/t$ as t increases, and is therefore too small to control instabilities in large times. This strategy therefore fails.

In this paper, we will directly prove that the series (2.16) converges as M goes to $+\infty$, in analytic spaces. This leads to a direct construction of a genuine solution of Navier Stokes equations, defined by

$$u^\nu(t, x, y) = U + \sum_{j=1}^{+\infty} \nu^{Nj} u^j(t, x, y) \quad (2.18)$$

The underlying idea is the following: if we try to control the difference between the true solution and an approximate one, we have to bound solutions of linearized Navier Stokes equations. However because of the shear, vertical derivatives of such solutions increase polynomially in time, simply because of the term $\partial_t + U(y)\partial_x$, which generates high normal derivatives. This polynomial growth can not be avoided, except if we are working with a finite sum of eigenmodes. For eigenmodes, we simply have an exponential growth, without polynomial disturbances. As a matter of fact, all the terms appearing in (2.18) are driven by eigenmodes through Orr Sommerfeld equations.

The proof of the convergence of (2.18) relies on the accurate description of the Green function of Orr Sommerfeld equations, detailed in [10], and on the introduction of so called generator functions. Generator functions combine all the norms of all the u^j , and can be seen as a time and space depending norm. We prove that these generator functions satisfy a Hopf inequality, which allows us to get analytic bounds which are uniform in M .

The plan of this paper is the following. We begin with the definition of generator functions. We then study the generator function of solutions of Laplace equations, and then of Orr Sommerfeld equations. We then detail the construction of u^j and derive uniform bounds on the generator functions, which ends the proof.

3 Generator functions

3.1 Definition

Let $f(x, y)$ be a smooth function. For $z_1, z_2 \geq 0$, we define the following two functions, called in this paper "generator functions"

$$\begin{aligned} Gen_0(f)(z_1, z_2) &= \sum_{\alpha \in \mathbb{Z}} \sum_{\ell \geq 0} e^{z_1|\alpha|} \|\partial_y^\ell f_\alpha\|_{\ell,0} \frac{z_2^\ell}{\ell!}, \\ Gen_\delta(f)(z_1, z_2) &= \sum_{\alpha \in \mathbb{Z}} \sum_{\ell \geq 0} e^{z_1|\alpha|} \|\partial_y^\ell f_\alpha\|_{\ell,\delta} \frac{z_2^\ell}{\ell!}, \end{aligned} \tag{3.1}$$

in which $f_\alpha(y)$ denotes the Fourier transform of $f(x, y)$ with respect to the x variable. In these sums,

$$\begin{aligned} \|f_\alpha\|_{\ell,0} &= \sup_y \varphi(y)^\ell |f_\alpha(y)|, \\ \|f_\alpha\|_{\ell,\delta} &= \sup_y \varphi(y)^\ell |f_\alpha(y)| \left(\delta^{-1} e^{-y/\delta} + 1 \right)^{-1}, \end{aligned}$$

where

$$\varphi(y) = \frac{y}{1+y}$$

and where the boundary layer thickness δ is equal to

$$\delta = \gamma_0 \nu^{1/4}$$

for some sufficiently large $\gamma_0 > 0$. More precisely, γ_0 will be chosen so that $\gamma_0^{-1} \leq \sqrt{\Re \lambda_0 / 2}$, where λ_0 is the maximal unstable eigenvalue of the linearized Euler equations around U .

Note that Gen_0 , Gen_δ and all their derivatives are non negative for positive z_1 and z_2 . These generator functions $Gen_0(\cdot)$ and $Gen_\delta(\cdot)$ will respectively control the velocity and the vorticity of the solutions of Navier Stokes equations.

For convenience, we introduce the following generator functions of one-dimensional functions $f = f(y)$:

$$\begin{aligned} Gen_{0,\alpha}(f)(z_2) &= \sum_{\ell \geq 0} \|\partial_y^\ell f\|_{\ell,0} \frac{z_2^\ell}{\ell!}, \\ Gen_{\delta,\alpha}(f)(z_2) &= \sum_{\ell \geq 0} \|\partial_y^\ell f\|_{\ell,\delta} \frac{z_2^\ell}{\ell!}. \end{aligned} \tag{3.2}$$

Of course, it follows that

$$Gen_0(f) = \sum_{\alpha \in \mathbb{Z}} e^{z_1|\alpha|} Gen_{0,\alpha}(f_\alpha)$$

for functions of two variables $f = f(x, y)$, and similarly for Gen_δ .

3.2 Properties

For any $\ell, \ell' \geq 0$, we have

$$\begin{aligned} \|f\|_{\ell,\delta} &\leq \|f\|_{\ell,0}, & \|f\|_{\ell+1,\delta} &\leq \|f\|_{\ell,\delta}, \\ \|fg\|_{\ell,\delta} &\leq \|f\|_{\ell',0} \|g\|_{\ell-\ell',\delta}. \end{aligned} \quad (3.3)$$

Next, we have the following Proposition

Proposition 3.1. *Let f and g be two functions. For non negative z_1 and z_2 , there hold*

$$\begin{aligned} Gen_\delta(fg) &\leq Gen_0(f)Gen_\delta(g), \\ Gen_\delta(\partial_x f) &= \partial_{z_1} Gen_\delta(f), & Gen_\delta(\partial_x^2 f) &= \partial_{z_1}^2 Gen_\delta(f), \\ Gen_\delta(\varphi \partial_y f) &\leq C_0 \partial_{z_2} Gen_\delta(f), \end{aligned}$$

for some universal constant C_0 , provided $|z_2|$ is small enough.

Proof. First, note that

$$(fg)_\alpha = \sum_{\alpha' \in \mathbb{Z}} f_{\alpha'} g_{\alpha-\alpha'},$$

and

$$\partial_y^\beta (fg)_\alpha = \sum_{\alpha' \in \mathbb{Z}} \sum_{0 \leq \beta' \leq \beta} \frac{\beta!}{\beta'!(\beta-\beta')!} \partial_y^{\beta'} f_{\alpha'} \partial_y^{\beta-\beta'} g_{\alpha-\alpha'}.$$

Thus,

$$\begin{aligned} &Gen_\delta(fg)(z_1, z_2) \\ &= \sum_{\alpha \in \mathbb{Z}} \sum_{\beta \geq 0} e^{z_1|\alpha|} \|\partial_y^\beta (fg)_\alpha\|_{\beta,\delta} \frac{z_2^\beta}{\beta!} \\ &\leq \sum_{\alpha \in \mathbb{Z}} \sum_{\beta \geq 0} \sum_{\alpha' \in \mathbb{Z}} \sum_{0 \leq \beta' \leq \beta} e^{z_1|\alpha|} \|\partial_y^{\beta'} f_{\alpha'}\|_{\beta',0} \|\partial_y^{\beta-\beta'} g_{\alpha-\alpha'}\|_{\beta-\beta',\delta} \frac{z_2^\beta}{\beta'!(\beta-\beta')!} \\ &\leq \sum_{\alpha, \alpha' \in \mathbb{Z}} \sum_{\beta \geq 0} \sum_{\beta \geq \beta'} e^{z_1|\alpha|} e^{z_1|\alpha-\alpha'|} \|\partial_y^{\beta'} f_{\alpha'}\|_{\beta',0} \|\partial_y^{\beta-\beta'} g_{\alpha-\alpha'}\|_{\beta-\beta',\delta} \frac{z_2^\beta z_2^{\beta-\beta'}}{\beta'!(\beta-\beta')!} \\ &\leq Gen_0(f)(z_1, z_2) Gen_\delta(g)(z_1, z_2). \end{aligned}$$

Next, we write

$$\begin{aligned} Gen_\delta(\partial_x f) &= \sum_{\alpha \in \mathbb{Z}} \sum_{\ell \geq 0} e^{z_1 |\alpha|} \|\alpha \partial_y^\ell f_\alpha\|_{\ell, \delta} \frac{z_2^\ell}{\ell!} \\ &= \partial_{z_1} \sum_{\alpha \in \mathbb{Z}} \sum_{\ell \geq 0} e^{z_1 |\alpha|} \|\partial_y^\ell f_\alpha\|_{\ell, \delta} \frac{z_2^\ell}{\ell!} = \partial_{z_1} Gen_\delta(f), \end{aligned}$$

and similarly for $Gen_\delta(\partial_x^2 f)$. Finally, we compute

$$\begin{aligned} Gen_\delta(\varphi \partial_y f) &= \sum_{\alpha \in \mathbb{Z}} \sum_{\ell \geq 0} e^{z_1 |\alpha|} \|\partial_y^\ell(\varphi \partial_y f_\alpha)\|_{\ell, \delta} \frac{z_2^\ell}{\ell!} \\ &\leq \sum_{\alpha \in \mathbb{Z}} \sum_{\ell \geq 0} \sum_{0 \leq \ell' \leq \ell} e^{z_1 |\alpha|} \|\partial_y^{\ell'} \varphi \partial_y^{\ell - \ell' + 1} f_\alpha\|_{\ell, \delta} \frac{z_2^\ell}{\ell'! (\ell - \ell')!} \\ &\leq \left(1 + \sum_{\ell' \geq 0} \|\partial_y^{\ell'} \varphi\|_{0,0} \frac{z_2^{\ell'}}{\ell'!}\right) \sum_{\alpha \in \mathbb{Z}} \sum_{\ell - \ell' \geq 0} e^{z_1 |\alpha|} \|\partial_y^{\ell - \ell' + 1} f_\alpha\|_{\ell - \ell' + 1, \delta} \frac{z_2^{\ell - \ell'}}{(\ell - \ell')!} \\ &\leq C_0 \partial_{z_2} Gen_\delta(f), \end{aligned}$$

where we distinguished the cases $\ell' = 0$ and $\ell' > 0$. As φ is analytic, $\sum_{\ell' \geq 0} \|\partial_y^{\ell'} \varphi\|_{0,0} z_2^{\ell'} / \ell'!$ converges provided z_2 is small enough. The Proposition follows. \square

3.3 Generator function and divergence free condition

Note that for any functions u and g , Proposition 3.1 yields

$$Gen_\delta(u \partial_x g) \leq Gen_0(u) \partial_{z_1} Gen_\delta(g). \quad (3.4)$$

This is not true for $Gen_\delta(v \partial_y g)$, due to the boundary layer weight. We will investigate $Gen_\delta(v \partial_y g)$ when (u, v) satisfies the divergence free condition, namely

$$\partial_x u + \partial_y v = 0.$$

Precisely, we will prove the following Proposition.

Proposition 3.2. *For $|z_2| \leq 1$, there holds*

$$Gen_\delta(v \partial_y g) \leq C \left(Gen_0(v) + \partial_{z_1} Gen_0(u) \right) \partial_{z_2} Gen_\delta(g).$$

This Proposition is linked to the deep structure of Navier Stokes equations, namely to the precise link between the transport operator and the incompressibility condition. Note that we "loose" one derivative: our bound involves $\partial_x u$.

Proof. We compute

$$Gen_\delta(v\partial_y g) = \sum_{\alpha \in \mathbb{Z}} \sum_{\beta \geq 0} e^{z_1|\alpha|} \|\partial_y^\beta(v\partial_y g)_\alpha\|_{\beta,\delta} \frac{z_2^\beta}{\beta!},$$

in which

$$\partial_y^\beta(v\partial_y g)_\alpha = \sum_{\alpha' \in \mathbb{Z}} \sum_{0 \leq \beta' \leq \beta} \frac{\beta!}{\beta'!(\beta - \beta')!} \partial_y^{\beta'} v_{\alpha'} \partial_y^{\beta - \beta' + 1} g_{\alpha - \alpha'}.$$

For $\beta' > 0$, using the divergence-free condition $\partial_y v_\alpha = -i\alpha u_\alpha$, we estimate

$$\|\partial_y^{\beta'} v_{\alpha'} \partial_y^{\beta - \beta' + 1} g_{\alpha - \alpha'}\|_{\beta,\delta} \leq \|\alpha' \partial_y^{\beta' - 1} u_{\alpha'}\|_{\beta' - 1, 0} \|\partial_y^{\beta - \beta' + 1} g_{\alpha - \alpha'}\|_{\beta - \beta' + 1, \delta}.$$

On the other hand, for $\beta' = 0$, we estimate

$$\|v_{\alpha'} \partial_y^{\beta + 1} g_{\alpha - \alpha'}\|_{\beta,\delta} \leq \|\varphi^{-1} v_{\alpha'}\|_{0,0} \|\partial_y^{\beta + 1} g_{\alpha - \alpha'}\|_{\beta + 1, \delta}.$$

We note that for $y \geq 1$, $\varphi(y) \geq 1/2$ and hence

$$\|\chi_{\{y \geq 1\}} \varphi^{-1} v_{\alpha'}\|_{0,0} \leq 2 \|v_{\alpha'}\|_{0,0}.$$

When $y \leq 1$, using again the divergence-free condition, we write

$$v_{\alpha'}(y) = -i\alpha' \int_0^y u_{\alpha'}(y') dy' = -i\alpha' y \int_0^1 u_{\alpha'}(x, \theta y) d\theta.$$

Therefore, $\varphi(y)^{-1} |v_{\alpha'}(y)| \leq \sup_y |\alpha' u_{\alpha'}(y)|$ for $y \leq 1$. This proves that

$$\|\varphi^{-1} v_{\alpha'}\|_{0,0} \leq 2 \|v_{\alpha'}\|_{0,0} + \|\alpha' u_{\alpha'}\|_{0,0}.$$

Combining these inequalities for any $\alpha \in \mathbb{Z}$ and $\beta \geq 0$, we obtain

$$\begin{aligned} \|\partial_y^\beta(v\partial_y g)_\alpha\|_{\beta,\delta} &\leq \sum_{\alpha' \in \mathbb{Z}} (2 \|v_{\alpha'}\|_{0,0} + \|\alpha' u_{\alpha'}\|_{0,0}) \|\partial_y^{\beta + 1} g_{\alpha - \alpha'}\|_{\beta + 1, \delta} \\ &+ \sum_{\alpha' \in \mathbb{Z}} \sum_{1 \leq \beta' \leq \beta} \|\alpha' \partial_y^{\beta' - 1} u_{\alpha'}\|_{\beta' - 1, 0} \|\partial_y^{\beta - \beta' + 1} g_{\alpha - \alpha'}\|_{\beta - \beta' + 1, \delta} \frac{\beta!}{\beta'!(\beta - \beta')!}. \end{aligned}$$

It remains to multiply by $e^{z_1|\alpha|}z_2^\beta/\beta!$ and to sum all the terms over α, α', β and β' . The second term in the right hand side is bounded by the product of

$$\sum_{\alpha} \sum_{\beta} e^{|\alpha|z_1} \|\alpha \partial_y^\beta u_\alpha\|_{\beta,0} \frac{z_2^{\beta+1}}{(\beta+1)!},$$

which is bounded by $Gen_0(\partial_x u)$ provided $|z_2| \leq 1$ and of

$$\sum_{\alpha} \sum_{\beta} e^{|\alpha|z_1} \|\partial_y^{\beta+1} g_\alpha\|_{\beta+1} \frac{z_2^\beta}{\beta!},$$

which equals $\partial_{z_2} Gen_\delta(g)$. The first term is similar, which ends the proof. \square

Let us now bound derivatives of the transport term $u\partial_x g + v\partial_y g$.

Proposition 3.3. *Let*

$$\mathcal{A} = \left(Id + \partial_{z_1} + \partial_{z_2} \right) Gen_\delta$$

and

$$\mathcal{B} = Gen_0(u) + Gen_0(v) + \partial_{z_1} Gen_0(u) + \mathcal{A}(g).$$

Then

$$\mathcal{A}(u\partial_x g + v\partial_y g) \leq C\mathcal{B}\partial_{z_1}\mathcal{B} + C\mathcal{B}\partial_{z_2}\mathcal{B}.$$

Note that all the terms in \mathcal{A} are non negative, since all the derivatives of generator functions are non negative.

Proof. Let us successively bound all the terms appearing in $\mathcal{A}(u\partial_x g + v\partial_y g)$. First, $Gen_\delta(u\partial_x g + v\partial_y g)$ has been bounded in (3.4) and in the previous proposition. Next we compute

$$\begin{aligned} \partial_{z_1} Gen_\delta(u\partial_x g) &= Gen_\delta(\partial_x(u\partial_x g)) = Gen_\delta(\partial_x u\partial_x g + u\partial_x^2 g) \\ &\leq \partial_{z_1} Gen_0(u)\partial_{z_1} Gen_\delta(g) + Gen_0(u)\partial_{z_1}^2 Gen_\delta(g). \end{aligned} \quad (3.5)$$

Moreover, using Proposition 3.3,

$$\begin{aligned} \partial_{z_1} Gen_\delta(v\partial_y g) &= Gen_\delta(\partial_x(v\partial_y g)) = Gen_\delta(\partial_x v\partial_y g + v\partial_y \partial_x g) \\ &\leq C(\partial_{z_1} Gen_0(v) + \partial_{z_1}^2 Gen_0(u))\partial_{z_2} Gen_\delta(g) \\ &\quad + C(Gen_0(v) + \partial_{z_1} Gen_0(u))\partial_{z_2} \partial_{z_1} Gen_\delta(g). \end{aligned}$$

Let us now bound the term $\partial_{z_2} \text{Gen}_\delta(v \partial_y g)$. Precisely, we have to bound

$$\begin{aligned} & \frac{z_2^n}{n!} \|\varphi^{n+1} \partial_y^{n+1} (v_{\alpha'} \partial_y g_{\alpha-\alpha'})\|_{0,\delta} = \frac{z_2^n}{n!} \|\varphi^{n+1} \partial_y^n (\partial_y v_{\alpha'} \partial_y g_{\alpha-\alpha'} + v_{\alpha'} \partial_y^2 g_{\alpha-\alpha'})\|_{0,\delta} \\ & \leq \sum_{0 \leq k \leq n} \frac{z_2^n}{k!(n-k)!} \|\varphi^{n+1} \partial_y^{k+1} v_{\alpha'} \partial_y^{n+1-k} g_{\alpha-\alpha'} + \varphi^{n+1} \partial_y^k v_{\alpha'} \partial_y^{n+2-k} g_{\alpha-\alpha'}\|_{0,\delta}. \end{aligned}$$

Let us split this sum in two. The first sum equals, using the divergence free condition,

$$\begin{aligned} & \sum_{0 \leq k \leq n} \frac{z_2^n}{k!(n-k)!} \|\varphi^k \partial_y^k \partial_x u_{\alpha'} \varphi^{n+1-k} \partial_y^{n+1-k} g_{\alpha-\alpha'}\|_{0,\delta} \\ & \leq \sum_{0 \leq k \leq n} \frac{z_2^k}{k!} \|\varphi^k \partial_y^k \partial_x u_{\alpha'}\|_{0,0} \frac{z_2^{n-k}}{(n-k)!} \|\varphi^{n+1-k} \partial_y^{n+1-k} g_{\alpha-\alpha'}\|_{0,\delta}. \end{aligned}$$

Multiplying by $e^{|\alpha|z_1}$ and summing over α and α' , the sum is bounded by

$$\text{Gen}_0(\partial_x u) \partial_{z_2} \text{Gen}_\delta(g) = \partial_{z_1} \text{Gen}_0(u) \partial_{z_2} \text{Gen}_\delta(g).$$

On the other hand, the second sum equals to

$$\sum_{0 \leq k \leq n} \frac{z_2^n}{k!(n-k)!} \|\varphi^{n+1} \partial_y^k v_{\alpha'} \partial_y^{n+2-k} g_{\alpha-\alpha'}\|_{0,\delta}. \quad (3.6)$$

We follow the proof of the previous Proposition. First, for $k > 0$, this sum equals to

$$\sum_{1 \leq k \leq n} \frac{z_2^n}{k!(n-k)!} \|\varphi^{k-1} \partial_y^{k-1} \partial_x u_{\alpha'} \varphi^{n+2-k} \partial_y^{n+2-k} g_{\alpha-\alpha'}\|_{0,\delta}.$$

Multiplying by $e^{|\alpha|z_1}$, the corresponding sum is bounded by

$$\partial_{z_1} \text{Gen}_0(u) \partial_{z_2}^2 \text{Gen}_\delta(g),$$

provided that $|z_2| \leq 1$. It remains to bound the term $k = 0$ in (3.6):

$$\frac{z_2^n}{n!} \|\varphi^{n+1} v_{\alpha'} \partial_y^{n+2} g_{\alpha-\alpha'}\|_{0,\delta} \leq \left(2\|v_{\alpha'}\|_{0,0} + \|\alpha' u_{\alpha'}\|_{0,0}\right) \frac{z_2^n}{n!} \|\varphi^{n+2} \partial_y^{n+2} g_{\alpha-\alpha'}\|_{0,\delta}.$$

Multiplying by $e^{|\alpha|z_1}$, the corresponding sum is bounded by

$$(\text{Gen}_0(v) + \partial_{z_1} \text{Gen}_0(u)) \partial_{z_2}^2 \text{Gen}_\delta(g).$$

This leads to

$$\begin{aligned} \partial_{z_2} \text{Gen}_\delta(v \partial_y g) & \leq \partial_{z_1} \text{Gen}_0(u) \partial_{z_2} \text{Gen}_\delta(g) \\ & \quad + C_0 (\text{Gen}_0(v) + \partial_{z_1} \text{Gen}_0(u)) \partial_{z_2}^2 \text{Gen}_\delta(g). \end{aligned}$$

The bound on $\partial_{z_2} \text{Gen}_\delta(u \partial_x g)$ is similar which ends the proof of this Proposition. \square

4 Laplace equations

In this section we focus on the classical Laplace equation which is much easier than Orr Sommerfeld equations. We will apply the same arguments on Orr Sommerfeld in the next section.

4.1 In one space dimension

We consider the classical one-dimensional Laplace equation

$$\Delta_\alpha \phi = \partial_y^2 \phi - \alpha^2 \phi = f \quad (4.1)$$

on the half line $y \geq 0$, with Dirichlet boundary condition $\phi(0) = 0$ and $\lim_{y \rightarrow +\infty} \phi(y) = 0$. We recall that $\|f\|_{0,0} = \sup_{y \geq 0} |f(y)|$. Let us first recall the following classical result:

Proposition 4.1. (*L^∞ bounds*).

Let ϕ solve the one-dimensional Laplacian problem (4.1), with Dirichlet boundary condition. There holds

$$\alpha^2 \|\phi\|_{0,0} + |\alpha| \|\partial_y \phi\|_{0,0} + \|\partial_y^2 \phi\|_{0,0} \leq C \|f\|_{0,0}, \quad (4.2)$$

where the constant C is independent of the integer $\alpha \neq 0$.

Note that (4.2) states that we gain two derivatives by inverting the Laplace operator: a control on the maximum of f gives a control on the maximum of the first two derivatives of ϕ .

Proof. We will only consider the case $\alpha > 0$, the opposite case being similar. The Green function of $\partial_y^2 - \alpha^2$ is

$$G(x, y) = -\frac{1}{2\alpha} \left(e^{-\alpha|x-y|} - e^{-\alpha|x+y|} \right)$$

and its absolute value is bounded by $\alpha^{-1} e^{-\alpha|x-y|}$. The solution ϕ of (4.1) is explicitly given by

$$\phi(y) = \int_0^\infty G(x, y) f(x) dx. \quad (4.3)$$

A direct bound leads to

$$|\phi(y)| \leq \alpha^{-1} \|f\|_{0,0} \int_0^\infty e^{-\alpha|x-y|} dx \leq C \alpha^{-2} \|f\|_{0,0}$$

in which the extra α^{-1} factor is due to the x -integration. Splitting the integral formula (4.3) in $x < y$ and $x > y$ and differentiating it, we get

$$\|\partial_y \phi\|_{0,0} \leq C\alpha^{-1}\|f\|_{0,0}.$$

We then use the equation to bound $\partial_y^2 \phi$, which ends the proof of (4.2). \square

Next, in the case when f has a boundary layer behavior, we obtain the following result:

Proposition 4.2. (*Boundary layers norms*)

Let ϕ solve the one-dimensional Laplacian problem (4.1) with Dirichlet boundary condition. Provided $|\delta\alpha^2| \leq 1$, there hold

$$\|\nabla_\alpha \phi\|_{0,0} \leq C\|f\|_{0,\delta} \quad (4.4)$$

and

$$|\alpha|^2 \|\phi\|_{0,0} + \|\partial_y^2 \phi\|_{0,\delta} \leq C\|f\|_{0,\delta} \quad (4.5)$$

where the constant C is independent of the integer α .

Note that in the case of boundary layer norms, we only gain "one" derivative in supremum norm, but the usual two derivatives in boundary layer norm.

Proof. Using (4.3), we estimate

$$\begin{aligned} |\phi(y)| &\leq \alpha^{-1}\|f\|_{0,\delta} \int_0^\infty e^{-\alpha|y-x|} \left(1 + \delta^{-1}e^{-x/\delta}\right) dx \\ &\leq \alpha^{-1}\|f\|_{0,\delta} \left(\alpha^{-1} + \delta^{-1} \int_0^\infty e^{-x/\delta} dx\right) \end{aligned}$$

which yields the claimed bound for $\alpha\phi$. The bound on $\partial_y \phi$ is obtained by differentiating (4.3).

Let us turn to (4.5). Note that $|\partial_x G(x, y)| \leq 1$. As $G(0, y) = 0$ this gives $|G(x, y)| \leq |x|$. Therefore

$$|G(x, y)| \leq \min(\alpha^{-1}e^{-\alpha|x-y|}, |x|),$$

and hence

$$\begin{aligned} |\phi(y)| &\leq \|f\|_{0,\delta} \int_0^\infty \min(|x|, \alpha^{-1}e^{-\alpha|x-y|}) \left(\delta^{-1}e^{-x/\delta} + 1\right) dx \\ &\leq C\|f\|_{0,\delta} \left(\delta + \alpha^{-2}\right) \end{aligned}$$

which gives the desired bound when $|\delta\alpha^2| \leq 1$. We then use the equation to get the bound on $\|\partial_y^2 \phi\|_{0,\delta}$. \square

4.2 Laplace equation and generator functions

In this section, we will study the generator functions, introduced in Section 3.1, of solutions to the Laplace equation $\Delta\phi = \omega$. In the sequel, it is important to keep in mind that, in the application to Prandtl boundary layer stability, ω will have a boundary layer behavior, namely will behave like $\delta^{-1}e^{-Cy/\delta}$, whereas the stream function ϕ will be bounded in the limit.

Proposition 4.3. *Let $\phi_\alpha = \Delta_\alpha^{-1}\omega_\alpha$ on \mathbb{R}_+ with the Dirichlet boundary condition $\phi_\alpha|_{y=0} = 0$. For $|\delta\alpha^2| \leq 1$, there are positive constants C_0, θ_0 so that*

$$\text{Gen}_{\delta,\alpha}(\nabla_\alpha^2\phi_\alpha) + \text{Gen}_{0,\alpha}(\nabla\phi_\alpha) \leq C_0\text{Gen}_{\delta,\alpha}(\omega_\alpha), \quad (4.6)$$

for all z_2 so that $|z_2| \leq \theta_0$.

Moreover if $\phi = \Delta^{-1}\omega$ and if $\omega_\alpha = 0$ for all α such that $|\delta\alpha^2| \geq 1$, then

$$\text{Gen}_0(\nabla\phi) \leq C\text{Gen}_\delta(\omega), \quad (4.7)$$

$$\partial_{z_1}\text{Gen}_0(\nabla\phi) \leq C\partial_{z_1}\text{Gen}_\delta(\omega), \quad (4.8)$$

$$\partial_{z_2}\text{Gen}_0(\nabla\phi) \leq C\partial_{z_2}\text{Gen}_\delta(\omega) + \text{Gen}_\delta(\omega). \quad (4.9)$$

Proof. For $n \geq 1$, from the elliptic equation $\Delta_\alpha\phi_\alpha = \omega_\alpha$, we compute

$$\Delta_\alpha(\varphi^n\partial_y^n\phi_\alpha) = \varphi^n\partial_y^n\omega_\alpha + 2\partial_y(\varphi^n)\partial_y^{n+1}\phi_\alpha + \partial_y^2(\varphi^n)\partial_y^n\phi_\alpha.$$

Note that

$$\partial_y(\varphi^n)\partial_y^{n+1}\phi_\alpha = n\varphi'\varphi^{n-1}\partial_y^{n+1}\phi_\alpha,$$

and hence the $\|\cdot\|_{0,\delta}$ norm of this term is bounded by $n\|\varphi^{n-1}\partial_y^{n+1}\phi_\alpha\|_{0,\delta}$.

Moreover,

$$\partial_y^2(\varphi^n)\partial_y^n\phi_\alpha = \left(n(n-1)\varphi'^2\varphi^{n-2} + n\varphi''\varphi^{n-1}\right)\partial_y^n\phi_\alpha$$

whose $\|\cdot\|_{0,\delta}$ norm is bounded by $n(n-1)\|\varphi^{n-2}\partial_y^n\phi_\alpha\|_{0,\delta}$. Using Proposition 4.2, we get

$$\begin{aligned} & |\alpha|^2\|\varphi^n\partial_y^n\phi_\alpha\|_{0,0} + \|\partial_y^2(\varphi^n\partial_y^n\phi_\alpha)\|_{0,\delta} + \|\nabla_\alpha(\varphi^n\partial_y^n\phi_\alpha)\|_{0,0} \\ & \leq C\|\varphi^n\partial_y^n\omega_\alpha\|_{0,\delta} + Cn\|\varphi^{n-1}\partial_y^{n+1}\phi_\alpha\|_{0,\delta} + Cn(n-1)\|\varphi^{n-2}\partial_y^n\phi_\alpha\|_{0,\delta}. \end{aligned}$$

Expanding the left hand side, we get

$$\begin{aligned} & |\alpha|^2\|\varphi^n\partial_y^n\phi_\alpha\|_{0,0} + \|\varphi^n\partial_y^{n+2}\phi_\alpha\|_{0,\delta} + \|\varphi^n\partial_y^n\nabla_\alpha\phi_\alpha\|_{0,0} \\ & \leq C_0\|\varphi^n\partial_y^n\omega_\alpha\|_{0,\delta} + C_0n\|\varphi^{n-1}\partial_y^{n+1}\phi_\alpha\|_{0,\delta} \\ & + C_0n(n-1)\|\varphi^{n-2}\partial_y^n\phi_\alpha\|_{0,\delta} + C_0n\|\varphi^{n-1}\partial_y^n\phi_\alpha\|_{0,0}. \end{aligned} \quad (4.10)$$

Let

$$A_n = |\alpha|^2 \|\varphi^n \partial_y^n \phi_\alpha\|_{0,0} + \|\varphi^n \partial_y^{n+2} \phi_\alpha\|_{0,\delta} + \|\varphi^n \partial_y^n \nabla_\alpha \phi_\alpha\|_{0,0}.$$

Multiplying by $z_2^n/n!$ and summing over n , we get

$$\begin{aligned} \sum_{n \geq 0} A_n \frac{z_2^n}{n!} &\leq C_0 \sum_{n \geq 0} \|\varphi^n \partial_y^n \omega_\alpha\|_{0,\delta} \frac{z_2^n}{n!} + C_0 \sum_{n \geq 1} A_{n-1} \frac{z_2^n}{(n-1)!} \\ &\quad + C_0 \sum_{n \geq 2} A_{n-2} \frac{z_2^n}{(n-2)!} \end{aligned}$$

which ends the proof of (4.6), provided $|z_2|$ is small enough.

Next (4.7) is a direct consequence of (4.6), just summing in α . If we multiply (4.6) by $|\alpha|$ before summing it, this gives (4.8). Now we multiply (4.10) by $z_2^{n-1}/(n-1)!$ instead of $z_2^n/n!$. This gives

$$\begin{aligned} \sum_{n \geq 1} A_n \frac{z_2^{n-1}}{(n-1)!} &\leq C_0 \sum_{n \geq 1} \|\varphi^n \partial_y^n \omega_\alpha\|_{0,\delta} \frac{z_2^{n-1}}{(n-1)!} + C_0 \sum_{n \geq 1} A_{n-1} n \frac{z_2^{n-1}}{(n-1)!} \\ &\quad + C_0 \sum_{n \geq 2} n(n-1) A_{n-2} \frac{z_2^{n-1}}{(n-1)!}. \end{aligned}$$

The terms in the right hand side may be absorbed by the left hand side provided z_2 is small enough, except $C_0 A_0$, which is bounded by $Gen_{\delta,\alpha}(\omega_\alpha)$. This ends the proof of the Proposition. \square

5 Orr-Sommerfeld equations

5.1 Introduction

In this section, we study the Orr-Sommerfeld equations

$$Orr_{\alpha,c}(\phi) := -\epsilon \Delta_\alpha^2 \phi + (U - c) \Delta_\alpha \phi - U'' \phi = f, \quad (5.1)$$

together with the boundary conditions

$$\phi|_{y=0} = 0, \quad \partial_y \phi|_{y=0} = 0, \quad (5.2)$$

and $\phi \rightarrow 0$ as $y \rightarrow +\infty$, with $\Delta_\alpha = \partial_y^2 - \alpha^2$. We shall focus on the case when $\alpha \neq 0$; the $\alpha = 0$ case will be treated in Section 5.5. The Orr-Sommerfeld problem is the resolvent problem of the linearized Navier-Stokes equations around a shear profile U , written in terms of the stream function ϕ . We shall study the generator functions of Orr-Sommerfeld solutions.

Throughout this paper, $|\Im c|$ will always be larger than $3\Re\lambda_0/2$, where λ_0 is the speed of growth of the linear instability. In particular, $\Im c$ will be bounded away from 0. Moreover we will restrict ourselves to $\alpha \ll \epsilon^{-1/3}$, or precisely

$$|\epsilon\alpha^3 \log \nu| \leq 1. \quad (5.3)$$

Let us first describe Orr Sommerfeld equations in an informal way. For small ϵ , Orr Sommerfeld equations are a viscous perturbation of Rayleigh equations

$$Ray_{\alpha,c}(\phi) := (U - c)\Delta_\alpha\phi - U''\phi = f, \quad (5.4)$$

with boundary conditions $\phi(0) = 0$ and $\lim_{y \rightarrow +\infty} \phi(y) = 0$. Note that the equation $Ray_{\alpha,c}(\phi) = 0$ may be rewritten as

$$(\partial_y^2 - \alpha^2)\phi - \frac{U''}{U - c}\phi = 0.$$

As $y \rightarrow +\infty$, it therefore simplifies into $\partial_y^2\phi - \alpha^2\phi = 0$. Hence this equation has two independent solutions $\phi_{s,\pm}$, with respective asymptotic behaviors $e^{\pm|\alpha|y}$. Note that c is an eigenvalue if and only if $\phi_{s,-}(0) = 0$. We have to bound these solutions uniformly in $|\alpha| \geq 1$ and c , with $|\Im c| \geq 3\Re\lambda_0/2$. As α goes to ∞ , $\phi_{s,\pm}$ converge to $e^{\pm|\alpha|y}$. Moreover, $U''/(U - c)$ is bounded since $|\Im c| \geq 3\Re\lambda_0/2$ is bounded away from 0, therefore this term may be handled as a regular perturbation.

When we add the viscous term $\epsilon\Delta_\alpha^2\phi$, these two solutions are slightly perturbed, but give birth to two independent solutions of Orr Sommerfeld with a "slow" behavior, which behave like $e^{\pm|\alpha|y}$. Two additional solutions, called $\phi_{f,\pm}$, appear, with a fast behavior. For these solutions the viscous term is no longer negligible and is of the same order as the Rayleigh one. At leading order, $\phi_{f,\pm}$ are solutions to

$$-\epsilon\partial_y^4\phi + (U - c + \alpha^2\epsilon)\partial_y^2\phi = 0. \quad (5.5)$$

Let

$$\mu_f(y) = \sqrt{\alpha^2 + \frac{U - c}{\epsilon}},$$

taking the positive real part. Then at first order $\phi_{f,\pm}$ behaves like $e^{\pm \int_0^y \mu_f(z) dz}$.

Let $G_{\alpha,c}(x, y)$ be the Green function of the Orr-Sommerfeld problem. This Green function may be decomposed in a "slow part" G_s and a "fast part" G_f , such that

$$G_{\alpha,c}(x, y) = G_s(x, y) + G_f(x, y).$$

We recall the following theorem, which is the main result of [10, Theorem 2.1].

Theorem 5.1. *Let α, c be arbitrary, so that $|\Im c|$ is bounded away from 0 and $|\alpha \Im c| > 3\Re \lambda_0/2$. Then, there are universal positive constants C_0, θ_0 so that*

$$|G_s(x, y)| \leq \frac{C_0}{\mu_s(1 + |\Im c|)} \left(e^{-\theta_0 \mu_s |x-y|} + e^{-\theta_0 \mu_s |x+y|} \right) \quad (5.6)$$

$$|G_f(x, y)| \leq \frac{C_0}{m_f(1 + |\Im c|)} \left(e^{-\theta_0 m_f |x-y|} + e^{-\theta_0 m_f |x+y|} \right) \quad (5.7)$$

for all $x, y \geq 0$, in which

$$\mu_s = |\alpha|, \quad m_f = \inf_y \Re \mu_f(y), \quad (5.8)$$

with

$$\mu_f(y) = \sqrt{\alpha^2 + \frac{U-c}{\epsilon}},$$

taking the positive real part. Similar bounds hold for derivatives, namely for $k \geq 0$ and $l \geq 0$ with $k+l \leq 3$,

$$|\partial_x^k \partial_y^l G_s(x, y)| \leq \frac{C_{k,l}}{\mu_s^{1-k-l}(1 + |\Im c|)} \left(e^{-\theta_0 \mu_s |x-y|} + e^{-\theta_0 \mu_s (|x+y|)} \right) \quad (5.9)$$

$$|\partial_x^k \partial_y^l G_f(x, y)| \leq \frac{C_{k,l}}{m_f^{1-k-l}(1 + |\Im c|)} \left(e^{-\theta_0 m_f |x-y|} + e^{-\theta_0 m_f |x+y|} \right) \quad (5.10)$$

In the sequel, $\Im c$ will always be bounded away from 0, but can be very large. This theorem is the main result of [10, Theorem 2.1]. However, for the sake of completeness we sketch in the following lines the computation of the Green function, at a formal level. The Green function $G_{\alpha,c}(x, y)$ is constructed through the representation

$$G_{\alpha,c}(x, y) = \begin{cases} \sum_{k=s,f} d_k(x) \phi_{k,-}(y) + \sum_{k=s,f} e_k(x) \phi_{k,-}(y), & y > x > 0 \\ \sum_{k=s,f} d_k(x) \phi_{k,-}(y) + \sum_{k=s,f} f_k(x) \phi_{k,+}(y), & 0 < y < x \end{cases}$$

where the first sum takes care of the boundary condition and the second one of the singularity of the Green function near $x = y$.

It remains to compute d_k, e_k and f_k , so that $G_{\alpha,c}$ is continuous, together with its first two derivatives, so that $\varepsilon \partial_y^3 G_{\alpha,c}$ has a unit jump at $y = x$ and

so that $G_{\alpha,c}$ satisfies Dirichlet boundary condition together with its first derivative. The main contribution in $\varepsilon\partial_y^3 G_{\alpha,c}$ comes from fast modes since $\mu_f \gg \mu_s$. In order to get a unit jump at $y = x$ we have to choose, at leading order,

$$e_f(x) \sim -\frac{\phi_{f,-}(x)^{-1}}{2\varepsilon\mu_f(x)^3}, \quad f_f(x) \sim -\frac{\phi_{f,+}(x)^{-1}}{2\varepsilon\mu_f(x)^3}$$

Note that

$$\frac{1}{\varepsilon\mu_f^2} = \frac{1}{\varepsilon\alpha^2 + U - c}$$

and is therefore bounded by $C/(1 + |\Im c|)$ for large $|\Im c|$. With this choice of e_f and f_f , at leading order, $G_{\alpha,c}$ and its second derivatives are equal at x^+ and x^- . To match the first derivative we use $\phi_s(y)$. Note that

$$\partial_y \left(e_f(x)\phi_{f,-}(y) \right) \sim \frac{1}{2\varepsilon\mu_f^2} = \frac{1}{2} \frac{1}{\varepsilon\alpha^2 + U - c}$$

The jump of the first derivative of fast modes is therefore bounded by $C/(1 + |\Im c|)^{-1}$ for large $|\Im c|$. This jump is compensated by the slow modes. As a consequence, as the jump in the first derivatives of $e^{\pm|\alpha|y}$ is $|\alpha|$, $e_s(x)$ and $f_s(x)$ are of order $1/|\alpha|(1 + |\Im c|)$ for large $|\Im c|$. The bounds on d_k can be obtained in a similar way.

5.2 Pseudoinverse

We will also be interested in the case when $Orr_{\alpha,c}$ is not invertible. In this case, $Im(Orr_{\alpha,c})$ is of codimension 1, and the equation $Orr_{\alpha,c}(\phi) = f$ may be solved only if $\langle f, \phi_{\alpha,c} \rangle_{L^2} = 0$, where $\phi_{\alpha,c}$ spans the orthogonal of $Im(Orr_{\alpha,c})$. In [10] we show that we can construct an inverse of $Orr_{\alpha,c}$ on $Im(Orr_{\alpha,c})$ through a kernel $G = G_s + G_f$ which satisfies the same bounds as in Theorem 5.1. Moreover, the eigenmode of $Orr_{\alpha,c}$ does not lie in $Im(Orr_{\alpha,c})$. More precisely

Theorem 5.1. [10] *Let α be fixed. Let c_0 be a simple eigenvalue of $Orr_{\alpha,c}$ with corresponding eigenmode ϕ_{α,c_0} . Then there exists a bounded family of linear forms l^ν and a family of pseudoinverse operators Orr^{-1} such that for any stream function ϕ ,*

$$Orr_{\alpha,c} \left(Orr^{-1}(\phi) \right) = \phi - l^\nu(\phi)\phi_{\alpha,c_0}.$$

Moreover, Orr^{-1} may be defined through a Green function $G = G_s + G_f$ which satisfies (5.6) and (5.7).

5.3 Bounds on solutions of Orr Sommerfeld equations

Proposition 5.2. (*L^∞ norms*)

Let ϕ solve the Orr-Sommerfeld problem (5.1)-(5.2). For $|\epsilon\alpha^3| \leq 1$, $|\alpha\Im c| > 3\Re\lambda_0/2$ and $|\Im c|$ bounded away from 0, there hold

$$|\alpha|^2 \|\phi\|_{0,0} + |\alpha| \|\nabla_\alpha \phi\|_{0,0} + \|\nabla_\alpha^2 \phi\|_{0,0} \leq \frac{C_0}{1 + |\Im c|} \|f\|_{0,0} \quad (5.11)$$

and

$$\|\sqrt{\epsilon} \nabla_\alpha^3 \phi\|_{0,0} + \|\epsilon \nabla_\alpha^4 \phi\|_{0,0} \leq C_0 \|f\|_{0,0}. \quad (5.12)$$

Equations (5.11) and (5.12) express a classical regularity result: Orr Sommerfeld equation is a small fourth order elliptic perturbation of a second order elliptic equation. Therefore we gain the full control on two derivatives of the solution, and partial controls on third and fourth derivatives, with prefactors $\sqrt{\epsilon}$ and ϵ .

Proof. By construction, the solution ϕ is of the form

$$\phi(y) = \int_0^\infty G_{\alpha,c}(x,y) f(x) dx. \quad (5.13)$$

Hence,

$$\begin{aligned} |\phi(y)| &\leq C_0 \int_0^\infty \left(\frac{e^{-\theta_0 \mu_s |x-y|}}{\mu_s (1 + |\Im c|)} + \frac{e^{-\theta_0 m_f |x-y|}}{m_f (1 + |\Im c|)} \right) f(x) dx \\ &+ C_0 \int_0^\infty \left(\frac{e^{-\theta_0 \mu_s |x+y|}}{\mu_s (1 + |\Im c|)} + \frac{e^{-\theta_0 m_f |x+y|}}{m_f (1 + |\Im c|)} \right) f(x) dx \\ &\leq C_0 \|f\|_{0,0} (\mu_s^{-2} + m_f^{-2}) (1 + |\Im c|)^{-1}. \end{aligned}$$

We recall that $\mu_s = |\alpha|$ and that $\epsilon \mu_f^2 = \epsilon \alpha^2 + (U - c)$. Hence,

$$\frac{\alpha^2}{\mu_f^2} = \frac{2}{1 + \alpha(U - c)/(\epsilon \alpha^3)},$$

which is bounded since $|\alpha\Im c| > \Re\lambda_0$ and $|\epsilon\alpha^3| \leq 1$. This proves that

$$\|\alpha^2 \phi\|_{0,0} \leq C_0 (1 + |\Im c|)^{-1} \|f\|_{0,0}.$$

To get the bounds on $\alpha \partial_y \phi$ and $\partial_y^2 \phi$, we differentiate (5.13) with respect to y , splitting the integral in $x < y$ and $x > y$, and fulfill similar computations.

Similarly, we compute

$$\begin{aligned}
|\partial_y^3 \phi(y)| &\leq C_0(1 + |\Im c|)^{-1} \int_0^\infty \left(\mu_s^2 e^{-\theta_0 \mu_s |x-y|} + m_f^2 e^{-\theta_0 m_f |x-y|} \right) f(x) dx \\
&\quad + C_0(1 + |\Im c|)^{-1} \int_0^\infty \left(\mu_s^2 e^{-\theta_0 \mu_s |x+y|} + m_f^2 e^{-\theta_0 m_f |x+y|} \right) f(x) dx \\
&\leq C_0(\mu_s + m_f)(1 + |\Im c|)^{-1} \|f\|_{0,0}.
\end{aligned}$$

As $\sqrt{|\varepsilon|}|\alpha| \leq 1$, $\sqrt{|\varepsilon|}\mu_s \leq C$ and

$$\sqrt{|\varepsilon|}\mu_f = |\sqrt{\varepsilon\alpha^2 + U - c}| \leq C(1 + |\Im c|),$$

which yields the estimate for $\sqrt{\varepsilon}\partial_y^3 \phi$. For $\varepsilon\partial_y^4 \phi$, we directly use the Orr-Sommerfeld equation $Orr_{\alpha,c}(\phi) = f$. \square

Proposition 5.3. (*Boundary layer norms*)

Let ϕ solve the Orr-Sommerfeld problem (5.1)-(5.2), with source f having a boundary layer behavior. For $|\varepsilon\alpha^3 \log \nu| \leq 1$, $|\alpha \Im c| > 3\Re\lambda_0/2$ and $|\Im c|$ bounded away from 0, there holds

$$(1 + |\Im c|) \left(\|\nabla_\alpha \phi\|_{0,0} + \|\nabla_\alpha^2 \phi\|_{0,\delta} \right) + |\varepsilon| \|\nabla_\alpha^4 \phi\|_{0,\delta} \leq C_0 \|f\|_{0,\delta}. \quad (5.14)$$

Proof. Let $\chi(y)$ be a non negative function which equals 1 for $0 \leq y \leq 1$ and 0 for $y > 1$. Let us split the forcing term f in its boundary layer term and in its "inner term"

$$f = f_b + f_i$$

with

$$f_b(y) = \chi\left(\frac{y}{\delta \log \delta^{-1}}\right) f(y).$$

Note that $\|f_i\|_{0,0} \leq C\|f\|_{0,\delta}$ and

$$|f_b(x)| \leq C\|f\|_{0,\delta} \delta^{-1} e^{-y/\delta}.$$

Let ϕ_b and ϕ_i be the solutions of $Orr(\phi_b) = f_b$ and $Orr(\phi_i) = f_i$. Note that ϕ_i satisfies (5.14), thanks to the previous Proposition.

It remains to bound ϕ_b . For this we split the Green function in its fast part G_f and its slow part G_s . For the fast part we have to bound $G_f \star f_b$, which is a convolution between an exponentially decreasing kernel and a exponentially decreasing source. It is therefore bounded by $C\|f\|_{0,\delta} \delta^{-1} e^{-y/\delta}$ provided $m_f > 2\delta^{-1}$, which is the case provided γ_0 is large enough.

Let us turn to the slow part G_s . Let us first assume that $G_s(0, y) = 0$ for any positive y . Then $\partial_y^2 G_s(0, y) = 0$ for any positive y . As

$$|\partial_y^2 \partial_x G_s(x, y)| \leq C \frac{\mu_s^2}{(1 + |\Im c|)}$$

we have

$$|\partial_y^2 G_s(x, y)| \leq \frac{C \mu_s^2 x}{1 + |\Im c|},$$

noting the x factor on the right hand side. By convolution between G_s and f_b we have

$$|\partial_y^2 \phi_b(y)| \leq C \frac{|\delta \alpha^2|}{1 + |\Im c|} \|f\|_{0, \delta},$$

which leads to the desired bound, taking into account that $|\delta \alpha^2| \leq C$, provided that $G_s(0, y) = 0$ for any positive y .

However, it is not the case that $G_s(0, y) = 0$ for $y > 0$, but we rather have

$$G(0, y) = G_f(0, y) + G_s(0, y) = 0.$$

Therefore $\partial_y^2 G_s(0, y) = -\partial_y^2 G_f(0, y)$. For $y \geq \frac{1}{\theta_0 m_f} \log m_f$, we get

$$|\partial_y^2 G_s(0, y)| \leq \frac{C_0 m_f}{1 + |\Im c|} e^{-\theta_0 m_f y} \leq \frac{C_0}{1 + |\Im c|}.$$

On the other hand, for $y \leq \frac{1}{\theta_0 m_f} \log m_f$, we use

$$|\partial_y^3 G_s(0, y)| \leq \frac{C_0 \mu_s^2}{1 + |\Im c|}$$

to get

$$|\partial_y^2 G_s(0, y)| \leq \frac{C_0}{1 + |\Im c|} + \frac{C_0 m_f^{-1} \mu_s^2 \log m_f}{1 + |\Im c|}$$

which is bounded by a constant divided by $(1 + |\Im c|)$, upon recalling $m_f > 2\delta^{-1}$ and using the assumption $\alpha^2 \sqrt{\nu} \log \frac{1}{\nu} \leq 1$. This ends the proof of the bound on $\partial_y^2 \phi_b$. The bounds on ϕ_b and $\partial_y \phi_b$ are similar. \square

5.4 Generator functions

In this section, we study the generator of solutions to the Orr-Sommerfeld problem.

Proposition 5.4. *Let ϕ solve the Orr-Sommerfeld problem (5.1)-(5.2), with source term f . For $|\epsilon\alpha^3 \log \nu| \leq 1$, $|\alpha\Im c| > 3\Re\lambda_0/2$ and for $|\Im c|$ bounded away from 0, there are positive constants C_0, θ_0 (independent on ϵ and α) so that*

$$\text{Gen}_{0,\alpha}(\nabla_\alpha \phi) + \text{Gen}_{\delta,\alpha}(\nabla_\alpha^2 \phi) \leq \frac{C_0}{1 + |\Im c|} \text{Gen}_{\delta,\alpha}(f), \quad (5.15)$$

for all z_2 so that $0 \leq z_2 \leq \theta_0$. Moreover, provided $f_\alpha = 0$ if $|\epsilon\alpha^3 \log \nu| \geq 1$,

$$\text{Gen}_0(u) + \text{Gen}_\delta(\omega) \leq \frac{C_0}{1 + |\Im c|} \text{Gen}_\delta(f), \quad (5.16)$$

$$\partial_{z_1} \text{Gen}_0(u) + \partial_{z_1} \text{Gen}_\delta(\omega) \leq \frac{C_0}{1 + |\Im c|} \partial_{z_1} \text{Gen}_\delta(f), \quad (5.17)$$

and

$$\partial_{z_2} \text{Gen}_0(u) + \partial_{z_2} \text{Gen}_\delta(\omega) \leq \frac{C_0}{1 + |\Im c|} \left[\partial_{z_2} \text{Gen}_\delta(f) + \text{Gen}_\delta(f) \right]. \quad (5.18)$$

Proof. We estimate each term in the generator functions. The term $n = 0$ is already treated in Proposition 5.3. For $n \geq 1$, we compute

$$\begin{aligned} \text{Orr}_{\alpha,c}(\varphi^n \partial_y^n \phi) &= \varphi^n \partial_y^n f - 3\epsilon \partial_y \varphi^n \partial_y^{n+3} \phi - 6\epsilon \partial_y^2 \varphi^n \partial_y^{n+2} \phi \\ &\quad - 3\epsilon \partial_y^3 \varphi^n \partial_y^{n+1} \phi - \epsilon \partial_y^4 \varphi^n \partial_y^n \phi \\ &\quad + 4\epsilon \alpha^2 \partial_y \varphi^n \partial_y^{n+1} \phi + 2\epsilon \alpha^2 \partial_y^2 \varphi^n \partial_y^n \phi \\ &\quad + (U - c) \partial_y^2 \varphi^n \partial_y^n \phi + 2(U - c) \partial_y \varphi^n \partial_y^{n+1} \phi \\ &\quad + \sum_{1 \leq k \leq n} \frac{n!}{k!(n-k)!} \varphi^n \left(\partial_y^k U \partial_y^{n-k} \Delta_\alpha \phi - \partial_y^k U'' \partial_y^{n-k} \phi \right). \end{aligned} \quad (5.19)$$

Let us estimate each term on the right. For convenience, we set

$$\mathcal{A}_n := \|\varphi^n \partial_y^n \nabla_\alpha \phi\|_{0,0} + \|\varphi^n \partial_y^n \nabla_\alpha^2 \phi\|_{0,\delta} + |\epsilon|(1 + |\Im c|)^{-1} \|\varphi^n \partial_y^n \nabla_\alpha^4 \phi\|_{0,\delta},$$

for $n \geq 0$, and $\mathcal{A}_n = 0$ for negative n . As $\varphi = y/(1+y)$, we compute

$$\begin{aligned} &\|3\epsilon \partial_y \varphi^n \partial_y^{n+3} \phi + 6\epsilon \partial_y^2 \varphi^n \partial_y^{n+2} \phi + 3\epsilon \partial_y^3 \varphi^n \partial_y^{n+1} \phi + \epsilon \partial_y^4 \varphi^n \partial_y^n \phi\|_{0,\delta} \\ &\leq C_0(1 + |\Im c|) \sum_{k=1}^4 \frac{n!}{(n-k)!} \mathcal{A}_{n-k}, \\ &\|4\epsilon \alpha^2 \partial_y \varphi^n \partial_y^{n+1} \phi + 2\epsilon \alpha^2 \partial_y^2 \varphi^n \partial_y^n \phi\|_{0,\delta} \leq C\epsilon \alpha^2 \left[n\mathcal{A}_{n-1} + n(n-1)\mathcal{A}_{n-2} \right], \end{aligned}$$

and

$$\begin{aligned}\|(U - c)\partial_y \varphi^n \partial_y^{n+1} \phi\|_{0,\delta} &\leq C_0(1 + |\Im c|)n\mathcal{A}_{n-1} \\ \|(U - c)\partial_y^2 \varphi^n \partial_y^n \phi\|_{0,\delta} &\leq C_0(1 + |\Im c|) \left[n\mathcal{A}_{n-1} + n(n-1)\mathcal{A}_{n-2} \right].\end{aligned}$$

Finally, we treat the summation in (5.19). Set

$$\mathcal{B}_n = \|\partial_y^n U\|_{0,0} + \|\partial_y^n U''\|_{0,0}.$$

We estimate

$$\begin{aligned}&\sum_{1 \leq k \leq n} \frac{n!}{k!(n-k)!} \|\varphi^n (\partial_y^k U \partial_y^{n-k} \Delta_\alpha \phi - \partial_y^k U'' \partial_y^{n-k} \phi)\|_{0,\delta} \\ &\leq C_0 \sum_{1 \leq k \leq n} \frac{n!}{k!(n-k)!} \left(\|\partial_y^k U\|_{0,0} \|\varphi^{n-k} \partial_y^{n-k} \Delta_\alpha \phi\|_{0,\delta} \right. \\ &\quad \left. + \|\partial_y^k U''\|_{0,0} \|\varphi^{n-k} \partial_y^{n-k} \phi\|_{0,\delta} \right) \\ &\leq C_0 \sum_{1 \leq k \leq n} \frac{n!}{k!(n-k)!} \mathcal{B}_k \mathcal{A}_{n-k}.\end{aligned}$$

Thus, applying Proposition 5.3 to (5.19), we obtain, using $|\varepsilon \alpha^2| \leq 1$,

$$\begin{aligned}&\|\nabla_\alpha(\varphi^n \partial_y^n \phi)\|_{0,0} + \|\nabla_\alpha^2(\varphi^n \partial_y^n \phi)\|_{0,\delta} + |\varepsilon|(1 + |\Im c|)^{-1} \|\nabla_\alpha^4(\varphi^n \partial_y^n \phi)\|_{0,\delta} \\ &\leq \frac{C_0}{1 + |\Im c|} \|\varphi^n \partial_y^n f\|_{0,\delta} + C_0 \sum_{k=1}^4 \frac{n!}{(n-k)!} \mathcal{A}_{n-k} + C_0 \sum_{1 \leq k \leq n} \frac{n!}{k!(n-k)!} \mathcal{B}_k \mathcal{A}_{n-k}.\end{aligned}$$

Expanding the left hand side, we thus have

$$\mathcal{A}_n \leq \frac{C_0}{1 + |\Im c|} \|\varphi^n \partial_y^n f\|_{0,\delta} + C_0 \sum_{k=1}^4 \frac{n!}{(n-k)!} \mathcal{A}_{n-k} + C_0 \sum_{1 \leq k \leq n} \frac{n!}{k!(n-k)!} \mathcal{B}_k \mathcal{A}_{n-k}$$

for all $n \geq 0$. Multiplying the above equation by $z_2^n/n!$ and summing up the result in $n \geq 0$, we obtain

$$\begin{aligned}\sum_{n \geq 0} \mathcal{A}_n \frac{z_2^n}{n!} &\leq \frac{C_0}{1 + |\Im c|} \text{Gen}_{\delta,\alpha}(f) + C_0 \sum_{n \geq 0} \sum_{k=1}^4 \frac{n!}{(n-k)!} \mathcal{A}_{n-k} \frac{z_2^n}{n!} \\ &\quad + C_0 \sum_{n \geq 0} \sum_{k=1}^n \frac{n!}{k!(n-k)!} \mathcal{B}_k \mathcal{A}_{n-k} \frac{z_2^n}{n!}.\end{aligned}\tag{5.20}$$

Since $|z_2| \leq 1$, we compute

$$\sum_{n \geq 0} \sum_{k=1}^4 \frac{n!}{(n-k)!} \mathcal{A}_{n-k} \frac{z_2^n}{n!} \leq 4z_2 \sum_{n \geq 0} \mathcal{A}_n \frac{z_2^n}{n!},$$

which can be absorbed into the left hand side of (5.20), for sufficiently small z_2 . Similarly,

$$\sum_{n \geq 0} \sum_{k=1}^n \frac{n!}{k!(n-k)!} \mathcal{B}_k \mathcal{A}_{n-k} \frac{z_2^n}{n!} \leq C_0 \sum_{n \geq 0} \mathcal{A}_n \frac{z_2^n}{n!} \sum_{n \geq 1} \mathcal{B}_n \frac{z_2^n}{n!}$$

which is again absorbed into the left of (5.20), upon using the assumption that U is analytic, and that the sum of \mathcal{B}_n begins on $n \geq 1$. This ends the proof of (5.15).

Summing (5.15) gives (5.17). Multiplying by $|\alpha|$ before summing (5.15) we get (5.18). Now multiplying by $z_1^{n-1}/(n-1)!$ instead of $\frac{z_1^n}{n!}$ we get

$$\begin{aligned} \sum_{n \geq 1} \mathcal{A}_n \frac{z_2^{n-1}}{(n-1)!} &\leq \frac{C_0}{1 + |\Im c|} \partial_{z_2} \text{Gen}_{\delta, \alpha}(f) + C_0 \sum_{n \geq 1} \sum_{k=1}^4 \frac{n!}{(n-k)!} \mathcal{A}_{n-k} \frac{z_2^{n-1}}{(n-1)!} \\ &\quad + C_0 \sum_{n \geq 1} \sum_{k=1}^n \frac{n!}{k!(n-k)!} \mathcal{B}_k \mathcal{A}_{n-k} \frac{z_2^n}{n!}. \end{aligned}$$

If z_2 is small enough, the right hand side may be absorbed in the left hand side, excepted the terms involving \mathcal{A}_0 , which are bounded by $\text{Gen}_{\delta}(f)$. This ends the proof of the Proposition. \square

5.5 The $\alpha = 0$ case

In this section, we treat the case when $\alpha = 0$. In this case, the resolvent equation of the linearized Navier-Stokes equations simply becomes the resolvent equation for the heat equation

$$(\lambda - \sqrt{\nu} \partial_y^2) u_0 = F_0, \quad (5.21)$$

with $u_0 = 0$ at $y = 0$, where u_0 denotes the Fourier mode of the first component of velocity u at $\alpha = 0$. We have the following simple proposition.

Proposition 5.5. *Let u_0 solve (5.21) and let $\omega_0 = \partial_y u_0$ be the corresponding vorticity. For $\Re \lambda > 3\Re \lambda_0/2$, we have*

$$\begin{aligned} \text{Gen}_{0,0}(u_0) &\leq \frac{C_0}{1 + \Re \lambda} \text{Gen}_{0,0}(F_0), \\ \text{Gen}_{\delta,0}(\omega_0) &\leq \frac{C_0}{1 + \Re \lambda} \left(\text{Gen}_{0,0}(F_0) + \text{Gen}_{\delta,0}(\partial_y F_0) \right), \end{aligned} \quad (5.22)$$

for all $z_2 \geq 0$.

Proof. The solution u_0 to (5.21) with the zero boundary condition satisfies

$$u_0(y) = \int_0^\infty G_0(y, z) F_0(z) dz$$

where $G_0(y, z)$ denotes the Green function for $(\lambda - \sqrt{\nu} \partial_y^2)$ with the Dirichlet boundary condition. In particular, we have

$$|G_0(y, z)| \leq \nu^{-1/4} |\lambda|^{-1/2} e^{-\nu^{-1/4} \Re \sqrt{\lambda} |y-z|}.$$

In particular, for $\Re \lambda > 3\Re \lambda_0/2$, we estimate

$$\begin{aligned} |u_0(y)| &\leq \int_0^\infty \nu^{-1/4} |\lambda|^{-1/2} e^{-\nu^{-1/4} \Re \sqrt{\lambda} |y-z|} |F_0(z)| dz \\ &\leq C_0 (1 + \Re \lambda)^{-1} \sup_y |F_0(y)|. \end{aligned}$$

The estimates for derivatives are obtained in the same way as done in the previous section for the Orr-Sommerfeld equations. The proposition follows. \square

6 Construction of the instability

6.1 Iterative construction

Let us now describe the iterative construction of u^n and ω^n and of the infinite series which defines the solution (2.18). We start with the most unstable eigenmode (ψ_0, α_0, c_0) to the Orr-Sommerfeld problem; namely, we start with a solution of

$$Orr_{\alpha_0, c_0}(\psi_0) = 0,$$

with the zero boundary conditions on $y = 0$, such that $\alpha_0 \Im c_0$ is maximum. In fact we just need to start from a mode such that $\alpha_0 \Im c_0$ is strictly larger than half of this maximum. Up to a change of sign we may assume that $\alpha_0 > 0$. Up to a rescaling we may also assume that $\alpha_0 = 1$.

This mode corresponds to a complex solution $\nabla^\perp(e^{i\alpha_0(x-c_0t)}\psi_0(y))$ of the linearized Navier Stokes equations. We have to take the real part of this solution in order to deal with real valued solutions. Note that $(\bar{\psi}_0, -\alpha_0, \bar{c}_0)$ is also an eigenmode. We therefore sum up the two unstable eigenmodes corresponding to α_0 and $-\alpha_0$ and define

$$\psi^1(t, x, y) = e^{i\alpha_0(x-\Re c_0t)+\alpha_0\Im c_0t}\psi_0(y) + e^{-i\alpha_0(x-\Re c_0t)+\alpha_0\Im c_0t}\bar{\psi}_0(y).$$

Let ψ_α^1 be the Fourier transform of ψ^1 in x variable. Then all the ψ_α^1 vanish, except two of them, namely $\alpha = \pm\alpha_0 = \pm 1$. We then iteratively solve the resolvent equation of the linearized Navier-Stokes problem (2.17) for $u^n = \nabla^\perp \psi^n$. In term of vorticity $\omega^n = \Delta \psi^n$, the problem reads

$$\partial_t \omega^n + U \partial_x \omega^n - U'' \partial_x \psi^n - \sqrt{\nu} \Delta \omega^n = - \sum_{1 \leq j \leq n-1} (u^j \cdot \nabla) \omega^{n-j}, \quad (6.1)$$

together with the zero boundary condition on $u^n = \nabla^\perp \psi^n$. Precisely, we search for ψ^n under the form

$$\psi^n = \sum_{|\alpha| \leq n} \psi_\alpha^n e^{i\alpha(x - \Re c_0 t)} e^{n \Im c_0 t}, \quad (6.2)$$

where the sum runs on all the α which are multiples of α_0 . This yields, for $n \geq 2$,

$$\text{Orr}_{\alpha,c}(\psi_\alpha^n) = \frac{1}{i\alpha} \sum_{\alpha'} \sum_{1 \leq j \leq n-1} (u_{\alpha-\alpha'}^j \cdot \nabla_{\alpha'}) \omega_{\alpha'}^{n-j}, \quad (6.3)$$

in which $u_\alpha^j = \nabla_\alpha^\perp \psi_\alpha^j$ and $\omega_\alpha^j = \Delta_\alpha \psi_\alpha^j$, together with the zero boundary conditions on ψ^n and $\partial_y \psi^n$, and in which

$$c = \Re c_0 + in \frac{\alpha_0}{\alpha} \Im c_0.$$

Note that a α^{-1} factor appears in front of the source term, since Orr Sommerfeld is obtained by taking the vorticity of Navier Stokes equations and dividing by α . Note also that all but a finite number of ψ_α^n vanish. Again the sum runs on all the α which are multiple of α_0 . Note also that $|\Im c| \geq |\Im c_0|$ and is thus bounded away from 0.

As Proposition 5.4 only holds for $|\alpha^3 \epsilon| \leq 1$ or equivalently $|\alpha| \leq \nu^{-1/4}$, we will only retain the $|\alpha| \leq \nu^{-1/4}$ in the construction of ψ^n and restrict (6.2) to

$$\psi^n = \sum_{|\alpha| \leq \nu^{-1/4}} \psi_\alpha^n e^{i\alpha(x - \Re c_0 t)} e^{n \Im c_0 t}.$$

This leads to the introduction of the force

$$f^n = \sum_{|\alpha| > \nu^{-1/4}} \sum_{\alpha'} \sum_{1 \leq j \leq n-1} (u_{\alpha-\alpha'}^j \cdot \nabla_{\alpha'}) \omega_{\alpha'}^{n-j}$$

that will be estimated below.

6.2 Bounds on ψ^n

We prove the following.

Proposition 6.1. *Introduce the iterative norm*

$$\begin{aligned} G^n &= \text{Gen}_0(u^n) + \text{Gen}_0(v^n) + \partial_{z_1} \text{Gen}_0(u^n) \\ &\quad + \text{Gen}_\delta(\omega^n) + \partial_{z_1} \text{Gen}_\delta(\omega^n) + \partial_{z_2} \text{Gen}_\delta(\omega^n), \end{aligned} \quad (6.4)$$

for $n \geq 1$. Then, $G^n(z_1, z_2)$ are well-defined for sufficiently small z_1, z_2 , and in addition, there exists some universal constant C_0 so that

$$G^n \leq \frac{C}{n} \sum_{1 \leq j \leq n-1} \left(G^j \partial_{z_1} G^{n-j} + G^j \partial_{z_2} G^{n-j} \right). \quad (6.5)$$

Note that the derivatives appearing in (6.4) are non negative.

Proof. Applying Proposition 5.4 to the Orr-Sommerfeld equation (6.3), using $|\alpha| \leq \nu^{-1/4}$, and summing over α , we get

$$\mathcal{A}(\omega^n) \leq \frac{C_0}{n} \mathcal{A} \left(\sum_{1 \leq j \leq n-1} (u^j \cdot \nabla) \omega^{n-j} \right). \quad (6.6)$$

Moreover, using Proposition 4.3

$$\text{Gen}_0(u^n) + \text{Gen}_0(v^n) \leq C \text{Gen}_\delta(\omega^n)$$

and

$$\partial_{z_1} \text{Gen}_0(u^n) \leq C \partial_{z_1} \text{Gen}_\delta(\omega^n).$$

Proposition 3.3 then gives the desired bound. \square

6.3 Bounds on the generator function

Theorem 6.1. *For $n \geq 1$, let $G^n(z_1, z_2)$ be defined as in (6.4). Then, the series*

$$G(\tau, z_1, z_2) = \sum_{n=1}^{+\infty} G^n(z_1, z_2) \tau^{n-1}$$

converges, for sufficiently small τ, z_1 , and z_2 .

Proof. For $N \geq 1$, let us introduce the partial sum

$$G_N(\tau, z_1, z_2) := \sum_{n=1}^N G^n(z_1, z_2) \tau^{n-1},$$

for $\tau, z_1, z_2 \geq 0$. Note that G_N is a polynomial in τ , and thus well-defined for all times $\tau \geq 0$. We also note that all the coefficients $G^n(z_1, z_2)$ are positive. In particular, $G_N(\tau, z_1, z_2)$ is positive, and so are all its time derivatives (when $z_1 > 0$ and $z_2 > 0$). Moreover, $G_N(\tau, z_1, z_2)$, and all its derivatives, are increasing in N . We also observe that, at $\tau = 0$,

$$G_N(0, z_1, z_2) = G^1(z_1, z_2),$$

for all $N \geq 1$, and hence,

$$G(0, z_1, z_2) = \lim_{N \rightarrow \infty} G_N(0, z_1, z_2) = G^1(z_1, z_2).$$

Next, multiplying (6.5) by τ^{n-2} and summing up the result, we obtain the following partial differential inequality

$$\partial_\tau G_N \leq C G_N \partial_{z_1} G_N + C G_N \partial_{z_2} G_N,$$

for all $N \geq 1$. Therefore the generator function satisfies an Hopf-type equation, or more precisely an Hopf inequality.

As G_N is increasing in z_1 and z_2 , we focus on the diagonal $z_1 = z_2$, and introduce

$$F_N(\tau, z) = G_N(\tau, \theta(\tau)z, \theta(\tau)z)$$

for $\tau, z \geq 0$, where $\theta(\cdot)$ will be chosen later, with $\theta(0) = 1$. It follows that F_N satisfies

$$\partial_\tau F_N \leq (2C F_N + \theta'(\tau)z) \partial_z F_N. \quad (6.7)$$

Note that F_N is increasing in N . At $\tau = 0$, $F_N(0, z) = G_N(0, z, z) = G^1(z, z)$, which is independent on N . Let $\rho > 0$ be small enough such that

$$M_0 = \sup_{0 \leq z \leq \rho} G^1(z, z)$$

is well defined. We now define $\theta(\tau)$ in such a way that

$$4C M_0 + \theta'(\tau)\rho < 0,$$

with $\theta(0) = 1$. For instance, we take

$$\theta(\tau) = 1 - 6C M_0 \rho^{-1} \tau.$$

We will work on a time interval where $\theta(\tau) \geq 1/2$, namely on $[0, T_0]$ where $T_0 = \rho/12CM_0$. Let T_N be the largest time $\leq T_0$ such that $F_N(\tau, z) \leq 2M_0$, for $0 \leq \tau \leq T_N$ and $0 \leq z \leq \rho$. Note that T_N exists and is strictly positive, since F_N is well defined for all the positive times, and continuous in time. It remains to prove that $\inf_{N \geq 1} T_N$ is positive.

Let us define the characteristics curves $X_N(\tau, z)$ by solving

$$\partial_\tau X_N(\tau, z) = -2CF_N(\tau, X_N(\tau, z)) - \theta'(\tau)X_N(\tau, z),$$

together with $X_N(0, z) = z$. Observe that the characteristics are outgoing at $z = 0$ and $z = \rho$. Therefore the characteristics completely fill $[0, T_N] \times [0, \rho]$. Let us now introduce

$$\tilde{F}_N(\tau, z) = F_N(\tau, X_N(\tau, z)).$$

It follows from (6.7) that

$$\partial_\tau \tilde{F}_N(\tau, z) = \partial_\tau F_N + \partial_\tau X_N \partial_z F_N \leq 0.$$

As a consequence,

$$\sup_{0 \leq z \leq \rho} F_N(\tau, z) \leq \sup_{0 \leq z \leq \rho} F_N(0, z) = \sup_{0 \leq z \leq \rho} G^1(z, z) = M_0.$$

Therefore $T_N \geq T_0$, for all $N \geq 1$, and F_N is bounded uniformly on $[0, T_0] \times [0, \rho]$. We can therefore take the limit $N \rightarrow +\infty$. This leads to the convergence of $G_N(\tau, z_1, z_1)$ as $N \rightarrow \infty$ for $0 \leq \tau \leq T_0$ and for $0 \leq z_1 \leq \rho$. Since $G_N(\tau, z_1, z_2)$ is increasing in z_1, z_2 , the convergence of $G_N(\tau, z_1, z_2)$ as $N \rightarrow \infty$ follows. \square

6.4 End of proof

It remains to bound the force term f^n . For this we note that the cut off occurs for $|\alpha| \geq \nu^{-1/4}$ where the corresponding modes are exponentially small. The force term is therefore exponentially small itself, and therefore arbitrary small in any Sobolev space.

Now $\sum_n \omega^n \tau^n$ converges for τ small enough. Let

$$\tau = \nu^N e^{\alpha_0 \Im c_0 t}.$$

Then as long as τ remains small, namely as long as t remains small than $C \log \nu^{-1}$ for some constant C , this series converges and defines a solution of the full incompressible Navier Stokes equations. Note that the series defining u and v also converges in the same way. This prove Theorem 1.2. Then Theorem 1.1 follows by a simple rescaling $(T, X, Z) = \sqrt{\nu}^{-1}(t, x, z)$.

7 Time dependent shear flow

We now turn to the case where the shear flow U_s depends on time. Let Ω_s be the corresponding vorticity. Note that U_s is a solution of

$$\partial_t U_s - \nu \partial_{yy} U_s = 0$$

and hence depends on t through $\sqrt{\nu t}$. We put this dependency in the notation $U_s(\sqrt{\nu t}, y)$. The perturbation satisfies

$$\begin{aligned} \partial_t \omega + (U_s(0) \cdot \nabla) \omega + (u \cdot \nabla) \Omega_s(0) - \sqrt{\nu} \Delta \omega \\ = -(u \cdot \nabla) \omega - Q_1(\sqrt{\nu t}) \omega - Q_2(\sqrt{\nu t}) u \end{aligned} \quad (7.1)$$

where

$$Q_1(\sqrt{\nu t}) \omega = (U_s(\sqrt{\nu t}) - U_s(0)) \partial_x \omega$$

and

$$Q_2(\sqrt{\nu t}) u = (u \cdot \nabla) (\Omega_s(\sqrt{\nu t}) - \Omega_s(0)).$$

Note that

$$U_s(\sqrt{\nu t}, y) = U_s(0) + \sum_{k=1}^M (\sqrt{\nu t})^k U_s^k(y) + O((\sqrt{\nu t})^{M+1}).$$

As we are interested in times of order $\log \nu$, and keeping in mind that $\partial_x \omega$ is always bounded by $\nu^{-1/4}$, we can put the $O()$ in the forcing term.

We will fulfill a perturbative analysis and look for solutions of (7.1) of the form

$$\psi(t, x, y) = \sum_{n \geq 1} \sum_{p \geq 0} \sum_{q \geq 0} \sum_{\alpha \in \mathbb{Z}} e^{n \Im c_0 t} t^p \sqrt{\nu}^q e^{i \alpha (x - \Re c_0 t)} \psi_\alpha^{n,p,q}(y). \quad (7.2)$$

In fact it is sufficient to bound q by some large integer M , since we allow a small forcing term.

Putting (7.2) in (7.1) we get

$$i \alpha \text{Orr}_{\alpha,c}(\psi_\alpha^{n,p,q}) + (p+1) \psi_\alpha^{n,p+1,q} = Q^{n,p,q} + L^{n,p,q} \quad (7.3)$$

where

$$Q^{n,p,q} = \sum_{\alpha'} \sum_{1 \leq j \leq n-1} \sum_{0 \leq k \leq p} \sum_{0 \leq l \leq q} (u_{\alpha-\alpha'}^{j,k,l} \cdot \nabla_{\alpha'}) \omega_{\alpha'}^{n-j,p-k,q-l}$$

and

$$L^{n,p,q} = i\alpha \sum_{k=1}^M U_s^k \omega_\alpha^{n,p-k,q-k} - i\alpha \sum_{k=1}^M \psi_\alpha^{n,p-k,q-k} \partial_y \Omega_s^k,$$

with the convention that a quantity vanishes if one of its indices is negative. Note that $Q^{n,p,q}$ only involves $\psi_{\alpha'}^{n',p',q'}$ with $n' < n$. Next, $L^{n,p,q}$ involves $\psi_{\alpha'}^{n',p',q'}$ with $n' \leq n$ and $p' < p$ and $q' < q$.

We will solve this equation by recurrence on the power of $\sqrt{\nu}$, namely on q . We begin with the leading order $q = 0$. All the $\psi_\alpha^{n,p,0}$ vanish, except when $p = 0$. System (7.2) then reduces to (6.3)

$$\psi_\alpha^{n,0,0} = \psi_\alpha^n$$

which are constructed in the previous section, up to any arbitrarily large n .

We then turn to $q = 1$. The first terms $\psi_\alpha^{n,0,0}$ create an "error term" L involving $\sqrt{\nu}^k t^k$ for $1 \leq k \leq M$. Let us first focus on the case $k = 1$. For $k = 1$, the corresponding $L^{n,1,1}$ term is

$$i\alpha \left(U_s^1 \omega_\alpha^{n,0,0} - \psi_\alpha^{n,0,0} \partial_y \Omega_s^1 \right).$$

For $n = 1$ and $\alpha = \pm 1$ we note that $Orr_{\alpha,c}$ is not invertible. This operator may also be non invertible for other values of α (in finite number). To simplify the discussion we assume that this does not occur (the general case is similar).

If $(n, \alpha) \neq (1, \pm 1)$, we take $\psi_\alpha^{n,p,1} = 0$ for $p > 1$. This leads to

$$i\alpha Orr_{\alpha,c}(\psi_\alpha^{n,1,1}) = \sum_{\alpha'} \sum_{1 \leq j \leq n-1} \sum_{k=0,1} (w_{\alpha-\alpha'}^{j,k,k} \cdot \nabla_{\alpha'}) \omega_{\alpha'}^{n-j,1-k,1-k} + L^{n,1,1} \quad (7.4)$$

which is a linearized version of (6.3).

For $(n, \alpha) = (1, \pm 1)$, we note that $Orr_{\alpha,c}$ is not invertible. Let A be the right hand side of (7.4). Thanks to Theorem 5.1 we define

$$\psi_{\pm 1}^{1,1,1} = Orr^{-1}(A),$$

and we use $\psi_{\pm 1}^{1,2,1}$ to handle the remainder

$$2\omega_{\pm 1}^{1,2,1} = l^\nu(A) \phi_{1,c_0}.$$

Now to bound $\psi_\alpha^{n,1,1}$ we introduce the corresponding generator function G^1 and proceed as in the previous section. This leads to the following inequality

$$\partial_t G^1 \leq G^0 \partial_{z_1} G^1 + G^0 \partial_{z_2} G^1 + G^1 \partial_{z_1} G^0 + G^1 \partial_{z_2} G^0 + C \partial_x G^0 + C G^0.$$

Using the same arguments as in the previous section, we obtain bound uniform bounds on G^1 . The recurrence can be continued using similar arguments.

References

- [1] R. Alexandre, Y.-G. Wang, C.-J. Xu, and T. Yang. Well-posedness of the Prandtl equation in Sobolev spaces. *J. Amer. Math. Soc.*, 28(3):745–784, 2015.
- [2] P. G. Drazin and W. H. Reid. *Hydrodynamic stability*. Cambridge Mathematical Library. Cambridge University Press, Cambridge, second edition, 2004. With a foreword by John Miles.
- [3] D. Gérard-Varet and E. Dormy. On the ill-posedness of the Prandtl equation. *J. Amer. Math. Soc.*, 23(2):591–609, 2010.
- [4] D. Gerard-Varet, Y. Maekawa, and N. Masmoudi. Gevrey stability of Prandtl expansions for 2D Navier-Stokes flows. *Duke Math. J.* 167 (13) (2018), 2531-2631.
- [5] D. Gerard-Varet and N. Masmoudi. Well-posedness for the Prandtl system without analyticity or monotonicity. *Ann. Sci. Éc. Norm. Supér. (4)*, 48(6):1273–1325, 2015.
- [6] D. Gérard-Varet and T. T. Nguyen. Remarks on the ill-posedness of the Prandtl equation. *Asymptot. Anal.*, 77(1-2):71–88, 2012.
- [7] E. Grenier. On the nonlinear instability of Euler and Prandtl equations. *Comm. Pure Appl. Math.*, 53(9):1067–1091, 2000.
- [8] E. Grenier, Y. Guo, and T. T. Nguyen. Spectral stability of Prandtl boundary layers: an overview. *Analysis (Berlin)*, 35(4):343–355, 2015.
- [9] E. Grenier, Y. Guo, and T. T. Nguyen. Spectral instability of characteristic boundary layer flows. *Duke Math. J.*, 165(16):3085–3146, 2016.
- [10] E. Grenier and T. T. Nguyen. Green function of Orr-Sommerfeld equations away from critical layers *SIAM J. Math. Anal.* 51 (2019), no. 2, 1279-1296.
- [11] E. Grenier and T. T. Nguyen. Sublayer of prandtl boundary layers. *Arch. Ration. Mech. Anal.*, 229 (2018), no. 3, 1139-1151.
- [12] E. Grenier and T. T. Nguyen. On nonlinear instability of Prandtl’s boundary layers: the case of Rayleigh’s stable shear flows. *arXiv:1706.01282.*, 2017.

- [13] E. Grenier and T. T. Nguyen. Sharp bounds on linear semigroup of Navier-Stokes with boundary layer norms. *arXiv:1703.00881*, 2017.
- [14] Y. Guo and T. T. Nguyen. A note on Prandtl boundary layers. *Comm. Pure Appl. Math.*, 64(10):1416–1438, 2011.
- [15] D. Iftimie and F. Sueur. Viscous boundary layers for the Navier-Stokes equations with the Navier slip conditions. *Arch. Ration. Mech. Anal.*, 199(1):145–175, 2011.
- [16] Y. Maekawa. On the inviscid limit problem of the vorticity equations for viscous incompressible flows in the half-plane. *Comm. Pure Appl. Math.*, 67(7):1045–1128, 2014.
- [17] N. Masmoudi and F. Rousset. Uniform regularity for the Navier-Stokes equation with Navier boundary condition. *Arch. Ration. Mech. Anal.*, 203(2):529–575, 2012.
- [18] N. Masmoudi and T. K. Wong. Local-in-time existence and uniqueness of solutions to the Prandtl equations by energy methods. *Comm. Pure Appl. Math.*, 68(10):1683–1741, 2015.
- [19] O. A. Oleinik and V. N. Samokhin. *Mathematical models in boundary layer theory*, volume 15 of *Applied Mathematics and Mathematical Computation*. Chapman & Hall/CRC, Boca Raton, FL, 1999.
- [20] M. Paddick. Stability and instability of Navier boundary layers. *Differential Integral Equations* 27 (2014), no. 9-10, 893-930.
- [21] L. Prandtl. *Über flüssigkeits-bewegung bei sehr kleiner reibung*. pages 484–491, 1904.
- [22] M. Sammartino and R. E. Caflisch. Zero viscosity limit for analytic solutions, of the Navier-Stokes equation on a half-space. I. Existence for Euler and Prandtl equations. *Comm. Math. Phys.*, 192(2):433–461, 1998.
- [23] M. Sammartino and R. E. Caflisch. Zero viscosity limit for analytic solutions of the Navier-Stokes equation on a half-space. II. Construction of the Navier-Stokes solution. *Comm. Math. Phys.*, 192(2):463–491, 1998.
- [24] H. Schlichting and K. Gersten. *Boundary-layer theory*. Springer-Verlag, Berlin, enlarged edition, 2000. With contributions by Egon Krause

and Herbert Oertel, Jr., Translated from the ninth German edition by
Katherine Mayes.