

## Practice Exam #1

Do not flip the page until told to do so.

Name: \_\_\_\_\_

Problem	Grade	Points Possible
1		5
2		5
3		5
4		15
5		15
6		15
Total		60

### Useful Equations

$x(t) = x_0 + v_{0x}t + \frac{1}{2}a_x t^2$ $y(t) = y_0 + v_{0y}t + \frac{1}{2}a_y t^2$ $v_x(t) = v_{0x} + a_x t$ $v_y(t) = v_{0y} + a_y t$ $v_{fx}^2 = v_{0x}^2 + 2a_x \Delta x$ $v_{fy}^2 = v_{0y}^2 + 2a_y \Delta y$ $a_c = \frac{v^2}{r}$ $\theta(t) = \theta_0 + \omega_0 t + \frac{1}{2}\alpha t^2$ $\omega(t) = \omega_0 + \alpha t$ $\omega^2 = \omega_0^2 + 2\alpha \Delta \theta$ $\sum_i \vec{\tau}_i = I\vec{\alpha} = \frac{d\vec{L}}{dt}$ $x(t) = A \cos(\omega t + \phi_0)$ $\omega = 2\pi f = 2\pi/T$ $v_{max} = A\omega$ $a_{max} = A\omega^2$ $v = \sqrt{F_T/\mu}$ $v = \lambda f$ $\omega_{spring} = \sqrt{k/m}$ $\omega_{pendulum} = \sqrt{g/L}$ $k = 2\pi/\lambda$	$\sum_i \vec{F}_i = m\vec{a} = \frac{d\vec{p}}{dt}$ $\vec{p} = m\vec{v}$ $F_{fr} = \mu_{s,k} F_N$ $K = \frac{1}{2}mv^2$ $K = \frac{1}{2}I\omega^2$ $U = mgy \text{ (gravity)}$ $U = \frac{1}{2}kx^2 \text{ (spring)}$ $a = R\alpha$ $v = R\omega$ $\vec{L} = I\vec{\omega}$ $\vec{L} = \vec{r} \times \vec{p}$ $I = \sum_i m_i R_i^2$ $\vec{P}_0 = \vec{P}_f$ $\vec{L}_0 = \vec{L}_f$ $\Sigma p_{0x} = \Sigma p_{fx}$ $\Sigma p_{0y} = \Sigma p_{fy}$
---	---

**Question 1:** If you are told an object has a non-zero constant acceleration, what can you say about the velocity of the object?

- (a) The object's velocity is increasing.
- (b) The object's velocity is decreasing.
- (c) The object's velocity isn't changing.
- (d) The object's velocity is changing.

**Question 2:** If you throw a ball up into the air, which of the following correctly describes the motion of the ball when it hits the peak of its trajectory?

- (a) The velocity will be zero, and the acceleration will be equal to gravity pointing down.
- (b) The velocity will be non-zero, and the acceleration will be equal to gravity pointing down.
- (c) The velocity and acceleration will both be non-zero and pointing down.
- (d) The velocity and acceleration will both be zero.
- (e) The velocity will be pointing down and the acceleration will be zero.

**Question 3:** If two vectors  $\vec{a}$  and  $\vec{b}$  obey the following relationships, describe them in words and with a picture.

$$\vec{a} + \vec{b} = \vec{c} \quad \text{and} \quad a + b = c. \tag{1}$$

**Question 4:** The length of the day on Earth is increasing at a (small) rate of 1.0 ms each century.

- (a) How much longer is the day after 20 centuries?
- (b) What is the percent change in the length of the day after 20 centuries?

**Question 5:** The cable supporting an elevator snaps when the empty elevator car is at rest at the top of a 120-m-tall building.

- (a) With what speed does the elevator strike the ground?
- (b) How long is it falling?
- (c) What is its speed half way down?
- (d) How long does it take to get half way down?

**Question 6:** For the following three vectors,

$$\begin{aligned}\vec{a} &= 4\hat{i} + 5\hat{j} - 6\hat{k} \\ \vec{b} &= -1\hat{i} + 2\hat{j} + 3\hat{k} \\ \vec{c} &= 4\hat{i} + 3\hat{j} + 2\hat{k},\end{aligned}$$

- (a) find  $\vec{r} = \vec{a} - \vec{b} + \vec{c}$ ;
- (b) find the angle between  $\vec{r}$  and the  $x$ -axis;
- (c) find the component of  $\vec{a}$  along the direction of  $\vec{b}$ .